

MULTISCALE MODELS OF TAXIS-DRIVEN PATTERNING IN BACTERIAL POPULATIONS

CHUAN XUE* AND HANS G. OTHMER†

Abstract. Spatially-distributed populations of various types of bacteria often display intricate spatial patterns that are thought to result from the cellular response to gradients of nutrients or other attractants. In the past decade a great deal has been learned about signal transduction, metabolism and movement in *E. coli* and other bacteria, but translating the individual-level behavior into population-level dynamics is still a challenging problem. However it is computationally impractical to use a strictly cell-based model to understand patterning in growing populations, since the total number of cells may reach $10^{12} - 10^{15}$ in some experiments. In the past phenomenological equations such as the Patlak-Keller-Segel equations have been used in modeling the cell movement that is involved in the formation of such patterns, but the question remains as to how the microscopic behavior can be correctly described by a macroscopic equation. Significant progress has been made for bacterial species that employ a ‘run-and-tumble’ strategy of movement, in that macroscopic equations based on simplified schemes for signal transduction and turning behavior have been derived [14, 13]. Here we extend previous work in a number of directions: (i) we relax a number of the assumptions on the attractant gradient made in previous derivations, (ii) we use a more general turning rate function that better describes the biological behavior, and (iii) we incorporate the effect of hydrodynamic forces that arise when cells swim in close proximity to a surface. We also develop a new approach to solving the moment equations derived from the transport equation to obtain macroscopic equations to any desired order. Numerical examples show that the solution of the lowest-order macroscopic equation agrees well with the solution obtained from a Monte Carlo simulation of cell movement under a variety of temporal protocols for the signal. We also apply the method to derive equations of chemotactic movement that are governed by multiple chemotactic signals.

Key words. chemotaxis equations, diffusion approximation, pattern formation, velocity-jump processes, transport equations

AMS subject classifications. 35Q80, 92B05

1. Introduction. New techniques in cell and molecular biology have produced huge advances in our understanding of signal transduction and cellular response in many systems, and this has led to better cell-level models for problems ranging from biofilm formation to embryonic development. However, many problems involve large numbers of cells ($\mathcal{O}(10^{12} - 10^{15})$), and detailed cell-based descriptions are computationally prohibitive at present. Thus rational techniques for incorporating cell-level knowledge into macroscopic equations are needed for these problems. One such problem arises when large numbers of individuals collectively organize into spatial patterns, as for instance in bacterial pattern formation and biofilms. In these systems the collective organization involves response to spatial gradients of attractants or repellents. When cells move toward (away from) favorable (unfavorable) conditions, the movement is called positive (negative) *taxis* if they adjust the direction of movement in response to the signal, and *kinesis* if the frequency of directional changes or the speed of movement is changed. If the active movement is in response to the gradient of a chemical we call it *chemotaxis* or *chemokinesis*. In this paper we focus on bacterial chemokinesis, which has been studied extensively in the bacterium *Escherichia coli*. Despite the clear difference in the type of response, both taxis and kinesis are lumped

*School of Mathematics, 204 Vincent Hall, University of Minnesota, Minneapolis, MN 55455 (cxue@math.umn.edu). Research supported by NIH grant GM 29123.

†School of Mathematics, 270A Vincent Hall, and also Digital Technology Center, University of Minnesota, Minneapolis, MN 55455 (othmer@math.umn.edu). Research supported by NIH grant GM 29123.

together in the literature, and we do not distinguish between them here.

Escherichia coli is a cylindrical enteric bacterium $\sim 1\text{--}2\text{ }\mu\text{m}$ long, that swims using a run-and-tumble strategy [4, 5, 37]. Each cell has 5-8 helical flagella that are several body lengths long, and each flagellum is rotated by a basal rotary motor embedded in the cell membrane. When all are rotated counterclockwise (CCW) the flagella form a bundle and propel the cell forward in a smooth ‘run’ at a speed $s \sim 10\text{--}30\text{ }\mu\text{m/s}$; when rotated clockwise (CW) the bundle flies apart, the cell stops essentially instantaneously because of its low Reynolds number, and it begins to “tumble” in place. After a random time the cell picks a new run direction with a slight bias in the direction of the previous run [6]. The alternation of runs and tumbles comprises the ‘run-and-tumble’ random movement of the cell. In the absence of a signal gradient the run and tumble times are exponentially distributed with means of 1 s and 0.1 s, respectively, but when exposed to a signal gradient, the run time is extended when the cell moves up (down) a chemoattractant (chemorepellent) gradient [6]. The molecular basis of signal transduction and motor control will be described in Section 2.

Under certain conditions, the collective population-level response to attractants produces intricate spatial patterns, even though each individual executes the simple ‘run-and-tumble’ strategy. For instance, in Adler’s capillary assay *E. coli* cells move up the gradient of a nutrient (an attractant), and the population forms moving bands or rings [1]. More recently, Budrene and Berg found that when *E. coli* moves up the gradient of a nutrient, they can also release another stronger chemoattractant. They studied the patterns in two experimental configurations, one in which a small inoculant of cells is introduced at the center of a semi-solid agar layer containing a single carbon source, such as succinate or other highly-oxidized intermediates of the TCA cycle. In this case the colony grows as it consumes the nutrients, cells secrete the chemoattractant aspartate, and a variety of spatial patterns of cell density develops during a two-day period, including outward-moving concentric rings, and symmetric arrays of spots and stripes. In the second type of experiment, wherein cells are grown in a thin layer of liquid medium with the same carbon source, a network-like pattern of high cell density forms from the uniform cell density, but this subsequently breaks into aggregates in 5-15 minutes. The formation of these patterns involves intercellular communication of millions of cells through the secreted chemoattractant aspartate, and thus detailed cell-based models of signal transduction, attractant release, and cell movement would be computationally expensive.

Heretofore, models of these and similar patterns have employed the classical Patlak-Keller-Segel (PKS) description of chemotactic movement [2, 34, 36, 35, 30]. Additional mechanisms assumed in these models include nonlinearity in the chemotactic coefficient, loss of motility under starvation conditions, or a second repellent or waste field. To understand the patterns formed in the soft agar, Brenner et al. [8] coupled the PKS chemotaxis equation with reaction-diffusion equations for both the attractant and nutrient, and proposed a minimal mechanism for the swarm ring and aggregate formation. They suggest that the motion of the swarm ring is driven by local nutrient depletion, with the integrity resulting from the high concentration of the attractant at the location of the ring; in contrast, the aggregates formed in the ring results from fluctuations near the unstable uniform cell density. However, the question of how to justify the chemotaxis equation from a microscopic description is not addressed in any of the foregoing models.

Recently significant progress has been made toward incorporating characteristics of the cell-level behavior into the classical description of chemotaxis [14, 13]. Using a

simplified description of signal transduction, these authors studied the parabolic limit of a velocity jump process that models the ‘run-and-tumble’ behavior of bacteria, and showed that the cell density n evolves according to the parabolic equation

$$(1.1) \quad \frac{\partial n}{\partial t} = \nabla \cdot \left(\frac{s^2}{N\lambda_0} \nabla n - \frac{bs^2 t_a G'(S)}{N\lambda_0(1+t_a\lambda_0)(1+t_e\lambda_0)} n \nabla S \right).$$

Here S is the attractant concentration; N is the space dimension, and s , λ_0 , b , t_e , t_a and $G(S)$ are parameters and functions that characterize the behavior of single cells. The authors assumed that (a) the signal function $G(S(\mathbf{x}))$ is time-independent, (b) the gradient of the signal is shallow as measured by $G'(S)\mathbf{v} \cdot \nabla S \sim \mathcal{O}(\varepsilon) \text{ sec}^{-1}$, (c) the turning rate depends linearly on the internal state of the cell ($\lambda = \lambda_0 - by_1$), and (d) the quasi-steady-state approximation is valid in estimating the higher order moments in the moment closure step. However, assumption (a) is often unrealistic in the processes of bacterial pattern formation, and assumption (c) imposes additional restrictions on y_1 , *i.e.*, $y_1 < \frac{\lambda_0}{b}$, in order to guarantee the positivity of the turning rate. Assumption (b) was used to justify the neglect of the higher order moments, and while analysis showed that (b) can alternatively be replaced by Assumption (d) in order to allow larger signal gradients, (b) is implicitly required in the perturbation analysis on the diffusion time and space scales, as will be shown later by an example.

In this paper we remove some of these restrictions. In section 3 we relax the assumptions (a) and (c) in order to allow time-dependent signals and a general turning frequency of the cells, and show that when (b) is violated, diffusion time and space scales are inapplicable. There we also develop a new method for solving the infinite system of the moment equations, which allows elimination of (d). The method involves systematic application of a solvability theorem to a perturbation expansion of the solution. In section 4 we compare the solution of the macroscopic chemotaxis equation and a stochastic simulation of chemotactic cell movement under a variety of temporal dynamics of the signal. In section 5 we extend the method to allow for external force terms in the transport equation. We illustrate the use of the resulting equation with an application to the model of spiral stream formation in *Proteus mirabilis* colonies [38], where a centripetal force is generated during the runs of the two-dimensional cell movement. Finally, we explore macroscopic chemotaxis equations for bacterial populations when exposed to several chemosignals in section 6. Before introducing the details of the analysis, we describe the cell-based model of bacterial pattern formation used in [38], which is based on a cartoon description of signal transduction introduced in [27].

2. The cell-based model. Bacterial cells are small, the swimmers we study here are typically 1-2 μm long. Therefore, we characterize their movement by their position $\mathbf{x} \in \mathbb{R}^N$ and velocity $\mathbf{v} \in V \subseteq \mathbb{R}^N$ as functions of time t . In the experiments of Budrene and Berg [9], the cell density is $\mathcal{O}(10^8) \text{ ml}^{-1}$, thus the average volume fraction of the cell population in the substrate is $\mathcal{O}(10^{-4})$. Even if in an aggregate cells are 100 times more crowded than average, the volume fraction would still be as small as $\mathcal{O}(10^{-2})$. Therefore, it is plausible to assume cells are well separated, so there is no mechanical interaction between them. This means that we can treat the movement of different cells as independent processes. In *E. coli* the cell speed is more or less constant throughout the movement, so we assume only the direction of the velocity changes during a tumble. In addition, since the mean tumbling time (~ 0.1 s) is much shorter than the run time (~ 1 s), we here neglect the tumbling time and assume that cells reorient immediately. In addition, we neglect the rotational

diffusion of cells during a run. Therefore, movement of cells can be characterized by independent velocity jump processes of the type introduced in [25] and later used in [17, 26, 14, 13]. The process is determined by a turning rate λ , and a turning kernel $T(\mathbf{v}, \mathbf{v}', \dots)$ which gives the probability density of turning from \mathbf{v}' to \mathbf{v} after making the decision to turn. Since T is a probability density it must satisfy

$$\int_V T(\mathbf{v}, \mathbf{v}', \dots) d\mathbf{v} = 1,$$

which means that no cells are lost during the reorientation. A generalization can be made to include the tumbling of cells as a separate resting phase [25]. In that case, the stochastic process would be determined by three parameters: the transition rate from the moving phase to the resting phase λ , the transition rate from the resting phase to the next moving phase denoted as μ , and the turning kernel T . It has been shown, in the absence of internal dynamics, that inclusion of a resting phase results in a re-scaling of the diffusion rate and the chemotactic sensitivity in the resulting macroscopic equation, which is essentially a re-scaling of time [26].

When there is no signal gradient, the turning rate λ is a constant, while in the presence of a signal gradient, λ depends on the current state of the flagella motor, which in turn is determined as the output of the underlying signal transduction network that transduces the extracellular signal into a change in rotational state.

Signal transduction in *E. coli* is a very complicated input-output process (Figure 2.1). Attractant binding to a receptor reduces the autokinase activity of the associated CheA, and therefore reduces the level of phosphorylated CheY_P, which is the output of the transduction network, on a fast time scale (~ 0.1 s). This constitutes the excitation component. Changes in the methylation level of the receptor by CheR and CheB restores the activity of the receptor complex to its pre-stimulus level on a slow time scale (seconds to minutes), which is called adaptation. Adaptation allows the cell to respond to further signals. The output CheY_P in turn changes the rotational bias of the flagella motors, and thus changes the run-and-tumble behavior [21, 37, 7].




FIG. 2.1. Signaling components and pathways for *E. coli* chemotaxis. Chemoreceptors (MCPs) span the cytoplasmic membrane (hatched lines), with a ligand-binding domain on the periplasmic side and a signaling domain on the cytoplasmic side. The cytoplasmic signaling proteins, denoted *Che* in the text, are identified by single letters, e.g., A = CheA. (From [33] with permission.)

Several detailed mathematical models have been proposed to model the entire signal transduction network [33, 32, 23, 31]. In the deterministic models, the state of

a cell can be described by a list of intracellular variables $\mathbf{y} = (y_1, y_2, \dots, y_q) \in \mathbb{R}^q$, and different models can be cast in the form of systems of ODEs


$$(2.1) \quad \frac{d\mathbf{y}}{dt} = f(S, \mathbf{y})$$

with different f , where $S(\mathbf{x}, t)$ is the signal concentration. In this article, we adopt a simplified cartoon description which is minimal yet captures the essential excitation and adaptation components:

$$(2.2) \quad \frac{dy_1}{dt} = \frac{G(S) - (y_1 + y_2)}{t_e},$$

$$(2.3) \quad \frac{dy_2}{dt} = \frac{G(S) - y_2}{t_a}.$$

Here t_e and t_a with $t_e \ll t_a$ are the excitation and adaptation time scales, \mathbf{x} is the current spatial position of the cell, and $G(S)$ a functional of the signal detected by the receptors. If we assume that there is no cooperative binding and the binding reaction



equilibrates rapidly, then

$$(2.4) \quad G(S) = G_0 \left(\frac{S}{K_D + S} \right)$$

with the binding coefficient $K_D = k^-/k^+$. $G(S)$ is bounded by G_0 since the receptors will be saturated at large concentrations of the attractant. The cartoon model has been shown to predict the input-output behavior of the full model [33] in response to step changes in the signal [14].

We may identify y_1 as the negative of the deviation of CheY_P from its steady state, and therefore, we assume the turning rate of each cell depends only on y_1 , *i.e.*,

$$\lambda = \lambda(y_1).$$

In addition, we assume that the turning kernel T is independent of \mathbf{x} and \mathbf{y} ,

$$T = T(\mathbf{v}, \mathbf{v}').$$

It has been shown experimentally that after a tumble, a cell has slight tendency to continue its previous direction of movement [4].

Finally, the above description of cell movement can be coupled with components of cell metabolism and cell division, and reaction-diffusion equations for the nutrient and attractant. We note that the description of cell movement used here comes directly from the biological observations; and by using reaction-diffusion equations for the chemicals, as in section 4, convection of the chemicals in the fluid flow is implicitly neglected. This approximation is valid here because the flow is very slow as a result of the small Reynolds number and low volume fraction of the cell population.

The Monte Carlo scheme can be used to simulate the model, but stochastic simulation can become extremely expensive because of cell division. Consider that cells double in 2 hours, and that the entire experimental process can last 2 days.

Assuming that n_0 cells are introduced into the petri dish initially, we would have $2^{48/2} \times n_0 \simeq 1.7 \times 10^7 n_0$ cells at the end; thus we need a higher level description. In the next section, we introduce a new method to embed the cell-level behavior in the population-level description, so as to derive an evolution equation of the cell density $n(\mathbf{x}, t)$ from the transport equation.

3. The transport equation and its diffusion limit absent external forces.

Let $p(\mathbf{x}, \mathbf{v}, \mathbf{y}, t)$ be the density of cells having position $\mathbf{x} \in \Omega \subset \mathbb{R}^N$, velocity $\mathbf{v} \in V \subset \mathbb{R}^N$, and internal states $\mathbf{y} \in \mathbb{R}^q$ at time $t \geq 0$, where V is a compact subset of \mathbb{R}^n and symmetric about the origin. Then the velocity jump process introduced above leads to the following transport equation when there is no cell growth

$$(3.1) \quad \frac{\partial p}{\partial t} + \nabla_{\mathbf{x}} \cdot (\mathbf{v}p) + \nabla_{\mathbf{y}} \cdot (\mathbf{f}p) = -\lambda(\mathbf{y})p + \int_V \lambda(\mathbf{y})T(\mathbf{v}, \mathbf{v}', \mathbf{y})p(\mathbf{x}, \mathbf{v}', \mathbf{y}, t) d\mathbf{v}'.$$

Here the left hand side of the equation describes the change of the population density due to the cell runs and the evolution of internal states, while the right hand side models the reorientation during the “tumbles”. A fundamental assumption in using a velocity jump process to model the “run-and-tumble” movement is that jumps occur instantaneously, and therefore the forces are Dirac functions. This approximation is appropriate for swimming bacteria since the Reynolds number is so small that cells stop instantly at the end of a run. A more general treatment can be found in [29].

In [26], a resting phase has been introduced to incorporate cell birth and death. While in some organisms it is true that cells stop to divide or give birth, the swimming bacterium *E. coli* has been observed to divide while swimming smoothly [3]. Thus the resting phase introduced is not necessary here. Therefore, by assuming that the growth rate r is a function of the local nutrient level $f(\mathbf{x}, t)$, the transport equation with cell growth reads

$$(3.2) \quad \frac{\partial p}{\partial t} + \nabla_{\mathbf{x}} \cdot (\mathbf{v}p) + \nabla_{\mathbf{y}} \cdot (\mathbf{f}p) = -\lambda(\mathbf{y})p + \int_V \lambda(\mathbf{y})T(\mathbf{v}, \mathbf{v}', \mathbf{y})p(\mathbf{x}, \mathbf{v}', \mathbf{y}, t) d\mathbf{v}' + r(f)p.$$

When cells grow in the exponential phase in a rich medium, r is a constant. By defining $p = \bar{p}e^{rt}$ and observing that \bar{p} satisfies equation (3.1), we can derive the equation for $\bar{n} = \int \bar{p} d\mathbf{x}$ and therefore $n = \bar{n}e^{rt}$. For this reason we begin with the transport equation (3.1) in the following derivation.

Define

$$\begin{aligned} z_1 &= y_1 \\ z_2 &= y_2 - G(S), \end{aligned}$$

then from the equations of y_1, y_2 , we obtain the system

$$(3.3) \quad \begin{aligned} \frac{dz_1}{dt} &= \frac{-z_1 - z_2}{t_e} \\ \frac{dz_2}{dt} &= -\frac{z_2}{t_a} - G'(S(x(t), t)) \left(\nabla S \cdot \mathbf{v} + \frac{\partial S}{\partial t} \right). \end{aligned}$$

The transport equation in the new internal variables (z_1, z_2) reads

$$(3.4) \quad \begin{aligned} \frac{\partial p}{\partial t} + \nabla_{\mathbf{x}} \cdot (\mathbf{v}p) + \frac{\partial}{\partial z_1} \left(\frac{-z_1 - z_2}{t_e} \right) + \frac{\partial}{\partial z_2} \left(-\frac{z_2}{t_a} - G'(S)(\nabla S \cdot \mathbf{v} + \frac{\partial S}{\partial t}) \right) p \\ = -\lambda(z_1)p + \lambda(z_1) \int_V T(\mathbf{v}, \mathbf{v}')p(\mathbf{v}') d\mathbf{v}'. \end{aligned}$$

This change of variables for the internal state makes the following analysis much simpler.

In the remainder of this section we relax a number of assumptions used in [14, 13] and present a new method to derive the chemotaxis equation in the diffusion limit of the transport equation (3.4). We first list the assumptions on the turning kernel and turning rate.

3.1. Assumptions on the turning kernel and turning rate. In our analysis we adopt the assumptions of the turning kernel T in [17, 13]. The notation used here coincides with that in [26, 13].

Define operator \mathcal{T} and its adjoint $\mathcal{T}^* : L^2(V) \rightarrow L^2(V)$ as follows:

$$(3.5) \quad (\mathcal{T}f)(\mathbf{v}) = \int_V T(\mathbf{v}, \mathbf{v}') f(\mathbf{v}') d\mathbf{v}', \quad (\mathcal{T}^*f)(\mathbf{v}) = \int_V T(\mathbf{v}', \mathbf{v}) f(\mathbf{v}') d\mathbf{v}'.$$

Denote by \mathcal{K} to be the non-negative cone of $L^2(V)$, $\mathcal{K} = \{f \in L^2(V) | f \geq 0\}$. The assumptions on the turning kernel $T \in L^2(V \times V)$ are

- A1: $T(\mathbf{v}, \mathbf{v}') \geq 0$, $\int_V T(\mathbf{v}, \mathbf{v}') d\mathbf{v} = \int_V T(\mathbf{v}', \mathbf{v}) d\mathbf{v} = 1$.
- A2: There are functions $u_0, \phi, \psi \in \mathcal{K}$ with the property $u_0 \neq 0$, $\phi > 0$ a.e., such that $u_0(\mathbf{v})\phi(\mathbf{v}') \leq T(\mathbf{v}, \mathbf{v}') \leq u_0(\mathbf{v})\psi(\mathbf{v}')$.
- A3: $\|\mathcal{T}\|_{\langle 1 \rangle^\perp} < 1$.

From these assumptions, one can prove [17] that (a) \mathcal{T} is a compact operator on $L^2(V)$, with spectral radius 1; (b) 1 is a simple eigenvalue with normalized eigenfunction $g(\mathbf{v}) \equiv 1$.

Next define the operators

$$(3.6) \quad \mathcal{A} = -I + \mathcal{T}, \quad \mathcal{A}^* = -I + \mathcal{T}^*.$$

Note that the operator \mathcal{L} defined in [26] is $\lambda\mathcal{A}$ here; in our derivation we use \mathcal{A} instead of \mathcal{L} because \mathcal{A} is independent of \mathbf{y} . One can easily prove that \mathcal{A} has the following properties:

- (i) $\|\mathcal{A}\| \leq 2$.
- (ii) $\mathcal{N}(\mathcal{A}) = \mathcal{N}(\mathcal{A}^*) = \langle 1 \rangle$,
 $\mathcal{R}(\mathcal{A}) = \mathcal{R}(\mathcal{A}^*) = \langle 1 \rangle^\perp = \{f \in L^2(V) | \int_V f(\mathbf{v}) d\mathbf{v} = 0\}$.
- (iii) $\forall \gamma$ with positive real part, $\gamma I - \mathcal{A}$ is invertible.

For the turning rate, we introduce a more general form than used in [14, 13]. We assume λ can be expanded to a Taylor series

$$\lambda = \lambda_0 - a_1 z_1 + a_2 z_1^2 - a_3 z_1^3 + \dots$$

with a radius of convergence at least $\max\{G_0, 1\}$, which implies that

$$(3.7) \quad \sum_{k=1}^{\infty} |a_k| < \infty.$$

The negative signs in the expansion of λ are introduced for the convenience of later analysis. The form $\lambda = \lambda_0 - by_1$ used in [14, 13] is a special case of this general form with $a_1 = b$ positive and $a_k = 0$ for all $k > 1$.

3.2. The parabolic scaling. To simplify the exposition we assume at first that excitation is much faster than other processes, that is, $t_e = 0$, $z_1 = -z_2$. The general result is simply stated later. Therefore the transport equation becomes

$$(3.8) \quad \frac{\partial p}{\partial t} + \nabla_x \cdot (\mathbf{v}p) + \frac{\partial}{\partial z_2} \left(-\frac{z_2}{t_a} - G'(S)(\nabla S \cdot \mathbf{v} + \frac{\partial S}{\partial t})p \right) \\ = (\lambda_0 + a_1 z_2 + a_2 z_2^2 + \dots)(-p + \int_V T(\mathbf{v}, \mathbf{v}')p(\mathbf{v}') d\mathbf{v}').$$

Since the total cell mass is conserved, we denote

$$(3.9) \quad N_0 = \int_{\Omega} \int_V \int_{\mathbb{R}} p dz_2 d\mathbf{v} d\mathbf{x},$$

and scale p by setting,

$$(3.10) \quad \hat{p} = \frac{p}{N_0}$$

The mean run time of *E. coli* is $T \simeq 1$ s, the speed is $10 \sim 30 \mu\text{m/s}$ [4], and a self-organized aggregate of cells has spatial dimension of $150 - 250 \mu\text{m}$ [24]. Thus, let $s_0 = 10 \mu\text{m/s}$, $L = 1$ mm, and re-scale the variables by setting,

$$\hat{\mathbf{v}} = \frac{\mathbf{v}}{s_0}, \quad \hat{x} = \frac{x}{L}, \quad \hat{t} = \frac{t}{T_p}, \quad \hat{V} = \frac{V}{s_0} \\ \hat{\lambda}_0 = \lambda_0 T, \quad \hat{a}_k = a_k T, \quad \hat{t}_a = \frac{t_a}{T}, \quad \epsilon = \frac{T s_0}{L} = 0.01, \quad T_p = \frac{T}{\epsilon^2}.$$

Therefore, $\hat{\mathbf{v}}, \hat{\mathbf{x}}, \hat{t}_a, \hat{\lambda}_0, \hat{a}_k \sim \mathcal{O}(1)$. We also re-scale

$$(3.11) \quad \hat{S} = \frac{S}{K_D}, \quad \hat{G}(\hat{S}) = G(\hat{S}K_D), \quad \hat{T}(\hat{\mathbf{v}}, \hat{\mathbf{v}}') = s_0^{N-1} T(\mathbf{v}, \mathbf{v}')$$

where K_D is the binding constant defined earlier.

In these variables equation (3.8) becomes, after dropping the hats,

$$(3.12) \quad \epsilon^2 \frac{\partial p}{\partial t} + \epsilon \nabla_x \cdot (\mathbf{v}p) + \frac{\partial}{\partial z_2} \left(-\frac{z_2}{t_a} - G'(S)(\epsilon \nabla S \cdot \mathbf{v} + \epsilon^2 \frac{\partial S}{\partial t})p \right) \\ = (\lambda_0 + a_1 z_2 + a_2 z_2^2 + \dots)(-p + \int_V T(\mathbf{v}, \mathbf{v}')p(\mathbf{v}') d\mathbf{v}').$$

Here the space and time variation of S enters at $\mathcal{O}(\epsilon)$ and $\mathcal{O}(\epsilon^2)$, respectively.

The goal of the moment closure method is to derive an approximating evolution equation for the cell density $n(\mathbf{x}, t)$ from the transport equation (3.12). To do that, ultimately we need to integrate (3.12) with respect to both z_2 and \mathbf{v} ¹. There are two places that one can apply the perturbation expansion: (a) to the incomplete moments, *viz.*, the z_2 -moments or \mathbf{v} -moments; (b) to the complete moments which are obtained by integrating both z_2 and \mathbf{v} . (b) will be used when there are external forces acted on the cells in section 5. However, in this section, we show that because the 0th-order z_2 -moment M_0^0 is independent of \mathbf{v} , applying the perturbation method to the z_2 -moments directly can lead to the approximating equation for $n(\mathbf{x}, t)$ with minimal assumptions.

¹In the case that the signal function depends on $n(\mathbf{x}, t)$, *i.e.*, $S = S(n, \mathbf{x}, t)$, we can approximate S by $S(n^0, \mathbf{x}, t)$, where n^0 is defined in the expansion $n = n^0 + \epsilon n^1 + \epsilon^2 n^2 + \dots$. This approximation introduces a term of $\mathcal{O}(\epsilon)$ into the transport equation (3.12), and thus won't change the equation derived later for n^0 .

3.3. The z_2 -moment equations. Define the moments of z_2 as follows:

$$(3.13) \quad M_j = \int z_2^j p dz_2, \quad \forall j = 0, 1, 2, 3, \dots \quad \mathbf{M} = (M_0, M_1, M_2, \dots)^t.$$

By multiplying equation (3.12) by 1, z_2^j/j for $j \geq 1$ and integrating, we obtain the moment equations in the following compact form:

$$(3.14) \quad \epsilon^2 \frac{\partial}{\partial t} \mathbf{\Lambda} \mathbf{M} + \epsilon \mathbf{v} \cdot \nabla_x \mathbf{\Lambda} \mathbf{M} = \epsilon^2 \mathbf{B} \mathbf{M} + \epsilon \mathbf{C} \mathbf{M} + \mathbf{D} \mathbf{M}.$$

Here

$$(3.15) \quad \mathbf{B} = -G'(S) \frac{\partial S}{\partial t} \mathbf{J}^t,$$

$$(3.16) \quad \mathbf{C} = -G'(S) (\nabla S \cdot \mathbf{v}) \mathbf{J}^t,$$

and

$$(3.17) \quad \mathbf{D} = -\frac{1}{t_a} \text{diag} \{0, 1, 1, \dots\} + \mathcal{A} \mathbf{\Lambda} (\lambda_0 \mathbf{I} + a_1 \mathbf{J} + a_2 \mathbf{J}^2 + \dots),$$

where \mathcal{A} is the operator defined in (3.6), $\mathbf{\Lambda} : l^\infty(L^2(V)) \rightarrow l^\infty(L^2(V))$ is a diagonal scaling operator $\mathbf{\Lambda} = \text{diag} \{1, 1, \frac{1}{2}, \frac{1}{3}, \dots\}$, and $\mathbf{J} : l^\infty(L^2(V)) \rightarrow l^\infty(L^2(V))$ is the shift operator that has ones on the upper diagonal entries:

$$(3.18) \quad \mathbf{J} = \begin{bmatrix} 0 & 1 & 0 & \dots \\ 0 & 0 & 1 & \dots \\ 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}.$$

One can easily prove that \mathbf{J} has the following properties:

$$(3.19) \quad \|\mathbf{J}\|_{l^\infty(L^2(V))} = 1, \quad \ker(\mathbf{J}) = \langle (1, 0, 0, 0, \dots)^t \rangle,$$

$$(3.20) \quad \{0\} \subset \ker(\mathbf{J}) \subset \ker(\mathbf{J}^2) \subset \dots \subset \ker(\mathbf{J}^k) \subset \dots \subset l^\infty(L^2(V)),$$

$$(3.21) \quad \bigcup_{k=1}^{\infty} \ker(\mathbf{J}^k) \subsetneq l^\infty(L^2(V)).$$

Therefore, \mathbf{B} and \mathbf{C} are bounded linear operators on $l^\infty(L^2(V))$. One can also easily prove that \mathbf{D} is a bounded linear operator on $l^\infty(L^2(V))$ under the assumptions on the turning kernel and turning rate introduced in section 3.1.

Since we are interested in the long-time dynamics, we will apply the regular perturbation method to solve the system (3.14). We explore two sets of assumptions. In the first, we assume that $\hat{G}'(\hat{S}) \frac{\partial \hat{S}}{\partial t}$ and $\hat{G}'(\hat{S}) \nabla \hat{S} \cdot \hat{\mathbf{v}}$ are of $\mathcal{O}(1)$ in the nondimensionalized variables, corresponding to $G'(S) \frac{\partial S}{\partial t} \sim \mathcal{O}(\epsilon^2) \text{sec}^{-1}$ and $G'(S) \nabla S \cdot \mathbf{v} \sim \mathcal{O}(\epsilon) \text{sec}^{-1}$ in the dimensional variables. We show in section 3.5 that this assumption leads to the same chemotaxis equation as in [13]. In the second, we relax the first set of assumptions to allow $\hat{G}'(\hat{S}) \frac{\partial \hat{S}}{\partial t}$ to be $\mathcal{O}(\frac{1}{\epsilon})$ in the nondimensionalized variables, or $G'(S) \frac{\partial S}{\partial t} \sim \mathcal{O}(\epsilon) \text{sec}^{-1}$ in the dimensional variables. This assumption means that a cell doesn't experience a significant change in the fraction of receptors bound during

an average run time. If the gradient is very large, this assumption may be violated and the characteristic space and time scale may be very different from those of the diffusion process. Therefore, the solution of the diffusion-limit chemotaxis equation will not be a good approximation of the underlying velocity jump process at the location where sharp spikes of the attractant arise. For this set of assumption, we show in section 3.6 that the equation for the first order approximation n^0 of the cell density remains the same, but the equation for higher order terms n^j depends on $\frac{\partial S}{\partial t}$. First however we prove a solvability theorem that will be used in the asymptotic analysis.

3.4. A solvability theorem. For $k \geq 1$, we introduce sub-matrix operators of \mathbf{D} defined by partitioning \mathbf{D} as follows

$$\mathbf{D} = \begin{bmatrix} \mathbf{E}_k & \mathbf{F}_k \\ 0 & \mathbf{G}_k \end{bmatrix}.$$

Here \mathbf{E}_k is the upper-left $k \times k$ submatrix of \mathbf{D} , \mathbf{F}_k is the upper-right $k \times \infty$ submatrix, and \mathbf{G}_k is the lower-right remainder. Written out,

$$\mathbf{E}_1 = [\lambda_0 \mathcal{A}], \quad \mathbf{F}_1 = \mathcal{A}[a_1, a_2, \dots],$$

and for $k > 1$,

$$\mathbf{E}_k = \begin{bmatrix} \lambda_0 \mathcal{A} & a_1 \mathcal{A} & \cdots & a_{k-1} \mathcal{A} \\ 0 & \lambda_0 \mathcal{A} - \frac{1}{t_a} & \cdots & a_{k-2} \mathcal{A} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & \frac{\lambda_0}{k-1} \mathcal{A} - \frac{1}{t_a} \end{bmatrix}, \quad \mathbf{F}_k = \mathcal{A} \begin{bmatrix} a_k & a_{k+1} & a_{k+2} & \cdots \\ a_{k-1} & a_k & a_{k+1} & \cdots \\ \cdots & \cdots & \cdots & \cdots \\ \frac{a_1}{k-1} & \frac{a_2}{k-1} & \frac{a_3}{k-1} & \cdots \end{bmatrix}.$$

for any $k \geq 1$,

$$\mathbf{G}_k = -\frac{1}{t_a} \mathbf{I} + \mathcal{A} \operatorname{diag} \left\{ \frac{1}{k}, \frac{1}{k+1}, \frac{1}{k+2}, \dots \right\} (\lambda_0 \mathbf{I} + a_1 \mathbf{J} + a_2 \mathbf{J}^2 + \cdots) = -\frac{1}{t_a} \mathbf{I} + \mathcal{A} \mathbf{\Lambda}_k \mathbf{\Phi},$$

with $\mathbf{\Lambda}_k \triangleq \operatorname{diag} \left\{ \frac{1}{k}, \frac{1}{k+1}, \frac{1}{k+2}, \dots \right\}$ and $\mathbf{\Phi} \triangleq \lambda_0 \mathbf{I} + a_1 \mathbf{J} + a_2 \mathbf{J}^2 + \cdots$.

Since components of \mathbf{D} are operators on the space $L^2(V)$, \mathbf{E}_k is an operator on $(L^2(V))^k$. Also by the assumption on the turning rate (3.7), $\mathbf{F}_k: l^\infty(L^2(V)) \rightarrow (L^2(V))^k$, $\mathbf{G}_k: l^\infty(L^2(V)) \rightarrow l^\infty(L^2(V))$. In the following theorem we prove that for any k , the operators \mathbf{G}_k are bounded and invertible. We denote the $l^\infty(L^2(V))$ norm by $\|\cdot\|$ and the corresponding operator norm by $|||\cdot|||$.

THEOREM 3.1. *For any $k \geq 1$, we have*

- (i) \mathbf{G}_k is bounded with $|||\mathbf{G}_k||| \leq \frac{1}{t_a} + \frac{1}{k} \|\mathcal{A}\|_{L^2(V)} (\lambda_0 + \sum_{j=1}^{\infty} |a_j|)$;
- (ii) \mathbf{G}_k is invertible, i.e., $\mathbf{G}_k \mathbf{W} = 0, \mathbf{W} \in l^\infty(L^2(V)) \implies \mathbf{W} = 0$.

Proof. (i) $\forall \mathbf{W} \in l^\infty(L^2(V))$, we have

$$\begin{aligned} \|\mathbf{\Phi} \mathbf{W}\| &= \max_{i=1}^{\infty} \|\lambda_0 W_i + \sum_{j=1}^{\infty} a_j W_{i+j}\| \leq \|\mathbf{W}\| \cdot (|\lambda_0| + \sum_{j=1}^{\infty} |a_j|), \\ \|\mathbf{\Lambda}_k \mathbf{W}\| &= \max_{i=1}^{\infty} \left| \frac{1}{k+i-1} W_i \right| \leq \frac{1}{k} \|\mathbf{W}\|, \\ \|\mathcal{A} \mathbf{W}\| &\leq \|\mathcal{A}\|_{L^2(V)} \cdot \|\mathbf{W}\|. \end{aligned}$$

Therefore, Φ , Λ_k and \mathcal{A} are bounded operators on $l^\infty(L^2(V))$. Since $\mathbf{G}_k = -\frac{1}{t_a}\mathbf{I} + \mathcal{A}\Lambda_k\Phi$, we have

$$|||\mathbf{G}_k||| \leq \frac{1}{t_a} + \|\mathcal{A}\|_{L^2(V)} |||\Lambda_k||| |||\Phi||| \leq \frac{1}{t_a} + \frac{1}{k} \|\mathcal{A}\|_{L^2(V)} (\lambda_0 + \sum_{j=1}^{\infty} |a_j|).$$

(ii) For $k > t_a \|\mathcal{A}\|_{L^2(V)} |||\Phi|||$, we have $|||t_a \mathcal{A}\Lambda_k\Phi||| < 1$. Therefore \mathbf{G}_k is invertible with $\mathbf{G}_k^{-1} = -\frac{1}{t_a} \sum_{i=0}^{\infty} (t_a \mathcal{A}\Lambda_k\Phi)^i$, *i.e.*, $\mathbf{G}_k \mathbf{W} = 0 \Rightarrow \mathbf{W} = 0$.

For $k \leq t_a \|\mathcal{A}\|_{L^2(V)} |||\Phi|||$, find $m > 0$ s.t. $k+m > t_a \|\mathcal{A}\|_{L^2(V)} |||\Phi|||$. Since \mathbf{G}_k is upper triangular, we get $W_j = 0, \forall j \geq m$ by observing \mathbf{G}_{k+m} is invertible; we then apply Gaussian elimination to the first $m-1$ equations in $\mathbf{G}_k \mathbf{W} = 0$ from the $(m-1)$ th row back to the 1st row to get $W_j = 0, j < m$. Property (iii) of the operator \mathcal{A} guarantees that Gaussian elimination applies. This completes the proof. \square

3.5. The regular perturbation of (3.14). Write \mathbf{M} as an expansion in powers of ϵ as

$$(3.22) \quad \mathbf{M} = \mathbf{M}^0 + \epsilon \mathbf{M}^1 + \epsilon^2 \mathbf{M}^2 + \dots$$

or

$$(3.23) \quad \begin{pmatrix} M_0 \\ M_1 \\ M_2 \\ M_3 \\ \vdots \end{pmatrix} = \begin{pmatrix} M_0^0 \\ M_1^0 \\ M_2^0 \\ M_3^0 \\ \vdots \end{pmatrix} + \epsilon \begin{pmatrix} M_0^1 \\ M_1^1 \\ M_2^1 \\ M_3^1 \\ \vdots \end{pmatrix} + \epsilon^2 \begin{pmatrix} M_0^2 \\ M_1^2 \\ M_2^2 \\ M_3^2 \\ \vdots \end{pmatrix} + \dots$$

The subscript indicates the order of the z_2 -moment, the superscript indicates the order of the term in the expansion.

After substituting (3.22) into the evolution equation (3.14) and comparing terms we find that

$\mathcal{O}(\epsilon^0)$:

$$\mathbf{D}\mathbf{M}^0 = 0,$$

By Theorem 3.1, we have

$$(3.24) \quad \mathcal{A}M_0^0 = 0 \quad \& \quad M_j^0 = 0, \forall j > 0.$$

By property (ii) of \mathcal{A} , we have M_0^0 independent of \mathbf{v} , *i.e.*, $M_0^0 = M_0^0(\mathbf{x}, t)$.

$\mathcal{O}(\epsilon^1)$:

$$\mathbf{v} \cdot \nabla_x \mathbf{A} \mathbf{M}^0 = \mathbf{C} \mathbf{M}^0 + \mathbf{D} \mathbf{M}^1,$$

or by using (3.24)

$$\begin{bmatrix} \mathbf{v} \cdot \nabla_x M_0^0 \\ G'(S) \nabla S \cdot \mathbf{v} M_0^0 \\ 0 \\ 0 \\ \vdots \end{bmatrix} = \mathbf{D} \mathbf{M}^1 = \begin{bmatrix} \mathbf{E}_2 & \mathbf{F}_2 \\ 0 & \mathbf{G}_2 \end{bmatrix} \mathbf{M}^1$$

Again, by Theorem 3.1, we have $M_j^1 = 0$, $\forall j > 1$, and the problem reduces to solving

$$\begin{aligned}\nabla_x \cdot \mathbf{v} M_0^0 &= \lambda_0 \mathcal{A} M_0^1 + a_1 \mathcal{A} M_1^1, \\ G'(S) \nabla S \cdot \mathbf{v} M_0^0 &= (\lambda_0 \mathcal{A} - \frac{1}{t_a}) M_1^1.\end{aligned}$$

By property (iii) of \mathcal{A} , $\lambda_0 \mathcal{A} - \frac{1}{t_a}$ is invertible, and thus,

$$\begin{aligned}M_1^1 &= (\lambda_0 \mathcal{A} - \frac{1}{t_a})^{-1} G'(S) \nabla S \cdot \mathbf{v} M_0^0, \\ \mathcal{A} M_0^1 &= \frac{1}{\lambda_0} \nabla_x \cdot \mathbf{v} M_0^0 - \frac{a_1 t_a}{\lambda_0} \mathcal{A} (t_a \lambda_0 \mathcal{A} - 1)^{-1} G'(S) \nabla S \cdot \mathbf{v} M_0^0.\end{aligned}$$

By property (ii) of \mathcal{A} , 0 is a simple eigenvalue, and we can define a pseudo-inverse operator of \mathcal{A} as $\mathcal{B} = (\mathcal{A}|_{\langle 1 \rangle^\perp})^{-1}$. Therefore, we obtain the representation,

$$(3.25) \quad M_0^1 = \mathcal{B} \frac{1}{\lambda_0} \mathbf{v} \cdot \nabla_x M_0^0 - \frac{a_1 t_a}{\lambda_0} (t_a \lambda_0 \mathcal{A} - 1)^{-1} G'(S) \nabla S \cdot \mathbf{v} M_0^0 + P_1,$$

where $P_1 \in \langle 1 \rangle$, *i.e.*, $P_1 = P_1(\mathbf{x}, t)$, is arbitrary. Notice that $n^1 = \int_V M_0^1 d\mathbf{v} = P_1 |V|$; thus n^1 can be determined once P_1 is known.

$\mathcal{O}(\epsilon^2)$:

$$\frac{\partial}{\partial t} \Lambda \mathbf{M}^0 + \mathbf{v} \cdot \nabla_x \Lambda \mathbf{M}^1 = \mathbf{B} \mathbf{M}^0 + \mathbf{C} \mathbf{M}^1 + \mathbf{D} \mathbf{M}^2$$

The first equation of the system implies

$$\frac{\partial}{\partial t} M_0^0 + \nabla_x \cdot \mathbf{v} M_0^1 \in \mathcal{R}(\mathcal{A}).$$

By property (ii) of \mathcal{A} ,

$$\int_V \left(\frac{\partial}{\partial t} M_0^0 + \mathbf{v} \cdot \nabla_x M_0^1 \right) d\mathbf{v} = 0.$$

Using (3.25), we get an equation for M_0^0

$$(3.26) \quad |V| \frac{\partial}{\partial t} M_0^0 + \frac{1}{\lambda_0} \nabla_x \cdot \int_V \mathbf{v} \mathcal{B} \mathbf{v} \cdot \nabla_x M_0^0 d\mathbf{v} - \frac{a_1 t_a}{\lambda_0} \nabla_x \cdot \int_V (\mathbf{v} (t_a \lambda_0 \mathcal{A} - 1)^{-1} G'(S) \nabla S \cdot \mathbf{v} M_0^0) d\mathbf{v} = 0.$$

By defining

$$(3.27) \quad D_n = -\frac{1}{|V| \lambda_0} \int_V \mathbf{v} \otimes \mathcal{B} \mathbf{v} d\mathbf{v}$$

and

$$(3.28) \quad \chi(S) = -\frac{a_1 t_a}{|V| \lambda_0} G'(S) \int_V \mathbf{v} \otimes (t_a \lambda_0 \mathcal{A} - 1)^{-1} \mathbf{v} d\mathbf{v},$$

we can rewrite equation (3.26) as

$$(3.29) \quad \frac{\partial}{\partial t} M_0^0 = \nabla_x \cdot (D_n \nabla_x M_0^0 - \chi(S) M_0^0 \nabla_x S).$$

The cell density $n(\mathbf{x}, t)$ is defined as

$$\begin{aligned} n &= \int_V \int_Z p(\mathbf{x}, \mathbf{v}, z_2, t) dz_2 d\mathbf{v} = \int_V M_0(\mathbf{x}, \mathbf{v}, t) d\mathbf{v} \\ &= \int_V (M_0^0 + \epsilon M_0^1 + \epsilon^2 M_0^2 + \dots) d\mathbf{v}, \end{aligned}$$

By expanding $n = n^0 + \epsilon n^1 + \epsilon^2 n^2 + \dots$, we find that

$$n^i = \int_V M_0^i d\mathbf{v}, \quad \forall i \geq 0.$$

In particular, $n^0 = |V| M_0^0$, thus $n = |V| M_0^0 + \mathcal{O}(\epsilon)$, and therefore we obtain the chemotaxis equation

$$(3.30) \quad \frac{\partial}{\partial t} n^0 = \nabla_x \cdot (D_n \nabla_x n^0 - \chi(S) n^0 \nabla_x S)$$

with a general tensor form of the diffusion rate (3.27) and the chemotaxis sensitivity (3.28).

Our standing assumption is that the cell speed is constant, and thus V is the a sphere of radius $s = \sqrt{\mathbf{v} \cdot \mathbf{v}}$ in 3-D. In the case that cells change direction of movement purely randomly, the turning kernel is given by the uniform density

$$(3.31) \quad T(\mathbf{v}, \mathbf{v}') = \frac{1}{|V|}.$$

In this case, the tensors D_n and $\chi(S)$ can be reduced to diagonal matrices, and thus scalars,

$$(3.32) \quad D_n = \frac{s^2}{N \lambda_0} I, \quad \chi(S) = G'(S) \frac{a_1 s^2 t_a}{N \lambda_0 (1 + t_a \lambda_0)}.$$

As a result, we obtain the classical chemotaxis equation for n^0

$$(3.33) \quad \frac{\partial}{\partial t} n^0 = \nabla_x \cdot \left(\frac{s^2}{N \lambda_0} \nabla_x n^0 - G'(S) \frac{a_1 s^2 t_a}{N \lambda_0 (1 + t_a \lambda_0)} n^0 \nabla_x S \right).$$

It is observed experimentally that the movement of *E. coli* shows directional persistence, and the turning kernel only depends on the angle θ between the old direction \mathbf{v}' and the new direction \mathbf{v} [6, 22], *i.e.*,

$$(3.34) \quad T(\mathbf{v}, \mathbf{v}') = h(\theta).$$

In this case, \mathcal{T} is a symmetric operator, the average velocity $\bar{\mathbf{v}}$ after reorientation

$$\bar{\mathbf{v}} = \int_V T(\mathbf{v}, \mathbf{v}') \mathbf{v} d\mathbf{v}$$

is parallel to the previous velocity \mathbf{v} , and thus the diffusion rate and the chemotaxis sensitivity are isotropic tensors (cf. [17], Theorem 3.5). As a result, one finds that $\mathcal{A}\mathbf{v} = -(1 - \psi_d)\mathbf{v}$ and

$$(3.35) \quad D_n = \frac{s^2}{N(1 - \psi_d)\lambda_0} I, \quad \chi(S) = G'(S) \frac{a_1 s^2 t_a}{N\lambda_0(1 + (1 - \psi_d)t_a\lambda_0)},$$

where

$$(3.36) \quad \psi_d = \frac{\bar{\mathbf{v}} \cdot \mathbf{v}'}{s^2} \in [-1, 1]$$

is the index of directional persistence introduced in [25]. We note that ψ_d can not be 1 in order to satisfy Assumption 2 on the turning kernel, and ψ_d has been reported to be about 0.33 in the wild-type *E. coli* [4]. From (3.35), we can see that the larger ψ_d is, the larger D_n and χ are, and therefore the larger the macroscopic chemotaxis velocity $\mathbf{u}_c = \chi(S)\nabla S$. The increase of \mathbf{u}_c to the persistence has also been analyzed in [20], where weak chemotaxis coupled with rotational diffusion was analyzed.

Equations for higher order terms. In order to obtain equations for higher order approximations of the cell density $n(\mathbf{x}, t)$, we can repeat the above calculation. The full equation system at $\mathcal{O}(\epsilon^2)$ is

$$\begin{bmatrix} \frac{\partial}{\partial t} M_0^0 + \mathbf{v} \cdot \nabla_x M_0^1 \\ \mathbf{v} \cdot \nabla_x M_1^1 + G'(S) \left(\frac{\partial S}{\partial t} M_0^0 + (\nabla S \cdot \mathbf{v}) M_0^1 \right) \\ G'(S) \nabla S \cdot \mathbf{v} M_1^1 \\ 0 \\ \vdots \end{bmatrix} = \begin{bmatrix} \lambda_0 \mathcal{A} M_0^2 + a_1 \mathcal{A} M_1^2 + a_2 \mathcal{A} M_2^2 + \cdots \\ (\lambda_0 \mathcal{A} - \frac{1}{t_a}) M_1^2 + a_1 \mathcal{A} M_2^2 + \cdots \\ (\frac{\lambda_0}{2} \mathcal{A} - \frac{1}{t_a}) M_2^2 + \cdots \\ (\frac{\lambda_0}{3} \mathcal{A} - \frac{1}{t_a}) M_3^2 + \cdots \\ \vdots \end{bmatrix}.$$

Similar to previous reasoning, we have $M_j^2 = 0, \forall j \geq 3$, and

$$\begin{aligned} M_2^2 &= 2(\lambda_0 \mathcal{A} - \frac{2}{t_a})^{-1} G'(S) (\nabla S \cdot \mathbf{v}) M_1^1, \\ M_1^2 &= (\lambda_0 \mathcal{A} - \frac{1}{t_a})^{-1} (\mathbf{v} \cdot \nabla_x M_1^1 + G'(S) \frac{\partial S}{\partial t} M_0^0 + G'(S) (\nabla S \cdot \mathbf{v}) M_0^1 - a_1 \mathcal{A} M_2^2), \\ M_0^2 &= \frac{\mathcal{B}}{\lambda_0} \mathbf{v} \cdot \nabla_x M_1^1 - \frac{a_1}{\lambda_0} M_1^2 - \frac{a_2}{\lambda_0} M_2^2 + P_2. \end{aligned}$$

Here, the term $(\mathcal{B}/\lambda_0)(\partial/\partial t M_0^0)$ in M_0^2 is absorbed into the \mathbf{v} -independent term P_2 . By considering the solvability condition of equations at the next order of ϵ , the equation for P_1 , and therefore, for $n^1 = P_1|V|$, can be obtained. Calculation reveals that the equation for n^1 is the same as n^0 in case that \mathbf{v} is an eigenfunction of \mathcal{T} , in particular for the turning kernel (3.34),

$$\frac{\partial}{\partial t} n^1 = \nabla_x \cdot \left(\frac{s^2}{N(1 - \psi_d)\lambda_0} \nabla_x n^1 - G'(S) \frac{a_1 s^2 t_a}{N\lambda_0(1 + (1 - \psi_d)t_a\lambda_0)} n^1 \nabla_x S \right).$$

If we force n^0 to satisfy the initial and boundary conditions of those for the cell density n , the higher order terms $n^j, j > 0$ should satisfy homogeneous initial and boundary conditions, and the zero mean constraint. Therefore, we conclude that $n^1 \equiv 0$, and thus, $n = n^0 + \mathcal{O}(\epsilon^2)$.

By allowing a finite excitation time in the cartoon model, we obtain the chemotaxis sensitivity tensor

$$(3.37) \quad \chi(S) = -\frac{a_1 t_a}{|V|\lambda_0} G'(S) \int_V \mathbf{v} \otimes (t_e \lambda_0 \mathcal{A} - 1)^{-1} (t_a \lambda_0 \mathcal{A} - 1)^{-1} \mathbf{v} d\mathbf{v},$$

as in [13], and using the turning kernel (3.34), the chemotaxis equation becomes

$$(3.38) \quad \frac{\partial}{\partial t} n^0 = \nabla \cdot \left(\frac{s^2}{N(1-\psi_d)\lambda_0} \nabla n^0 - \frac{a_1 s^2 t_a G'(S)}{N\lambda_0(1+(1-\psi_d)t_a\lambda_0)(1+(1-\psi_d)t_e\lambda_0)} n^0 \nabla S \right).$$

From this equation we can see that: (a) directional persistence increases both the diffusion rate and the macroscopic chemotactic velocity, as analyzed in [20]; (b) inclusion of the non-instantaneous excitation results in re-scaled chemotaxis sensitivity. The only difference by using the full cartoon model is, that instead of using matrix representations of \mathbf{M} and operators \mathbf{B} , \mathbf{C} , \mathbf{D} , block matrices should be used. A similar version of Theorem 3.1 can be proved without difficulty. One can also show that inclusion of a resting phase due to tumbling would result in a re-scaled diffusion rate and chemotaxis sensitivity by the fraction of running time.

3.6. A weaker assumption on the extracellular signal. In the above derivation we assumed that $G'(S) \frac{\partial S}{\partial t} \sim \mathcal{O}(1)$ on the parabolic (diffusion) time scale. However, when cells contribute to the signal field by secretion (example 4.2), $G'(S) \frac{\partial S}{\partial t}$ can become large when the cell density is large. Here we relax the assumption to allow $G'(S) \frac{\partial S}{\partial t} \sim \mathcal{O}(\frac{1}{\epsilon})$ on the parabolic time scale, which is $\mathcal{O}(\epsilon) \text{ sec}^{-1}$ in the dimensional variables. Under this assumption, we need to regroup the terms in the z_2 -moment equation (3.14). We define $\bar{S}_t = \epsilon \frac{\partial \hat{S}}{\partial t} \sim \mathcal{O}(1)$, $\bar{\mathbf{B}} = \epsilon \mathbf{B} \sim \mathcal{O}(1)$, then equation (3.14) can be rewritten as

$$(3.39) \quad \epsilon^2 \frac{\partial}{\partial t} \mathbf{A} \mathbf{M} + \epsilon \mathbf{v} \cdot \nabla_x \mathbf{A} \mathbf{M} = \epsilon (\bar{\mathbf{B}} + \mathbf{C}) \mathbf{M} + \mathbf{D} \mathbf{M}.$$

In this case, the equations in $\mathcal{O}(\epsilon)$ are

$$\mathbf{v} \cdot \nabla_x \mathbf{A} \mathbf{M}^0 = (\bar{\mathbf{B}} + \mathbf{C}) \mathbf{M}^0 + \mathbf{D} \mathbf{M}^1,$$

from which we can solve

$$(3.40) \quad M_1^1 = (\lambda_0 \mathcal{A} - \frac{1}{t_a})^{-1} G'(S) (\bar{S}_t + \nabla S \cdot \mathbf{v}) M_0^0,$$

$$(3.41) \quad M_0^1 = \mathcal{B} \frac{1}{\lambda_0} \mathbf{v} \cdot \nabla_x M_0^0 - \frac{a_1 t_a}{\lambda_0} (t_a \lambda_0 \mathcal{A} - 1)^{-1} G'(S) \nabla S \cdot \mathbf{v} M_0^0 + P_1.$$

In the representation of M_0^1 (3.41), the term $(t_a \lambda_0 \mathcal{A} - 1)^{-1} G'(S) \bar{S}_t M_0^0$ is absorbed by P_1 , since it is independent of \mathbf{v} . Therefore the equation for n_0 remains the same, *i.e.*, (3.30).

However, if we continue calculation for higher order terms, we obtain

$$\begin{aligned} M_2^2 &= 2(\lambda_0 \mathcal{A} - \frac{2}{t_a})^{-1} G'(S) (\bar{S}_t + \nabla S \cdot \mathbf{v}) M_1^1, \\ M_1^2 &= (\lambda_0 \mathcal{A} - \frac{1}{t_a})^{-1} (\mathbf{v} \cdot \nabla_x M_1^1 + G'(S) (\bar{S}_t + \nabla S \cdot \mathbf{v}) M_0^1 - a_1 \mathcal{A} M_2^2), \\ M_0^2 &= \frac{\mathcal{B}}{\lambda_0} \mathbf{v} \cdot \nabla_x M_0^1 - \frac{a_1}{\lambda_0} M_1^2 - \frac{a_2}{\lambda_0} M_2^2 + P_2. \end{aligned}$$

Here $a_1, a_2, \nabla S, \bar{S}_t, n^0$ and ∇n^0 enter the expression of M_0^2 , and by considering the solvability condition at $\mathcal{O}(\epsilon^3)$,

$$\int_V \frac{\partial}{\partial t} M_0^1 + \mathbf{v} \cdot \nabla_x M_0^2 d\mathbf{v} = 0,$$

we obtain an equation for n^1 ,

$$\begin{aligned} \frac{\partial}{\partial t} n^1 = \nabla_x \cdot \left[\frac{s^2}{N\lambda_0(1-\psi_d)} \nabla_x n^1 - G'(S) \frac{a_1 s^2 t_a}{N\lambda_0(1+t_a\lambda_0(1-\psi_d))} n^1 \nabla_x S \right] \\ + f(a_1, a_2, \nabla S, \bar{S}_t, n^0, \nabla n^0, \dots). \end{aligned}$$

The first-order term n^0 enters into the equation for n^1 through the function f which is linear in n^0 . In particular, for the turning kernel (3.34), f has the form

$$\begin{aligned} f = \nabla \cdot \left[\frac{a_1 t_a^2 s^2 \nabla(G'(S) \bar{S}_t n^0)}{N\lambda_0(1+\lambda_0 t_a(1-\psi_d))^2} \right. \\ + \frac{a_1 t_a s^2 G'(S) \bar{S}_t}{N\lambda_0(1+t_a\lambda_0(1-\psi_d))} \left(\frac{\nabla n^0}{\lambda_0(1-\psi_d)} - \frac{a_1 t_a n^0 G'(S) \nabla S}{\lambda_0(1+t_a\lambda_0(1-\psi_d))} \right) \\ \left. + \left(\frac{a_1^2 t_a(1-\psi_d)}{\lambda_0(1+\lambda_0 t_a(1-\psi_d))} + \frac{a_2}{\lambda_0} \right) \frac{4t_a^2 s^2 n^0 G'(S)^2 \bar{S}_t \nabla S}{N(2+\lambda_0 t_a(1-\psi_d))(1+\lambda_0 t_a(1-\psi_d))} \right]. \end{aligned}$$

In this case, the solution of the n^1 -equation is generally nonzero, and therefore $n = n^0 + \epsilon n^1 + \mathcal{O}(\epsilon^2)$, in contrast with the previous case.

4. Numerical comparison. According to the above perturbation analysis, the bacterial cell-based model in section 2 can be approximated by the solution of the chemotaxis equation (3.38) when coupled with an equation for the signal. In this section we first present two examples in 1-D to illustrate how accurate the approximation is. In both examples, we assume no cell growth and fast excitation, *i.e.*, $t_e = 0$; thus the equations for the internal dynamics become

$$(4.1) \quad \frac{dy_2}{dt} = \frac{G(S(\mathbf{x}, t)) - y_2}{t_a},$$

$$(4.2) \quad y_1 = G(S) - y_2.$$

with $G(S)$ defined by (2.4). We also assume no turning persistence ($\psi_d = 0$), and the turning rate

$$(4.3) \quad \lambda = \lambda_0 - \frac{2\lambda_0}{\pi} \tan^{-1}\left(\frac{y_1 \pi b}{2\lambda_0}\right),$$

which has the Taylor expansion,

$$\lambda = \lambda_0 - b y_1 + \dots$$

In this case, we compare with the stochastic simulation with the solution of

$$(4.4) \quad \frac{\partial}{\partial t} n = \nabla_x \cdot \left(\frac{s^2}{N\lambda_0} \nabla_x n - G'(S) \frac{b s^2 t_a}{N\lambda_0(1+t_a\lambda_0)} n \nabla_x S \right).$$

We then apply the 2-D version of both the continuum model and the cell-based model to the network-aggregate formation in *E. coli* colonies in section 4.3. The numerical method used in implementing the cell-based model is described in detail in Appendix A.

4.1. Aggregation and dispersion in one space dimension. In this example we analyze the motion of a bacterial population in response to a diffusing attractant on a periodic domain $[0, 4]$ mm. The dynamics of the attractant are described by the diffusion equation

$$(4.5) \quad \frac{\partial S}{\partial t} = D_s \Delta S,$$

with the initial condition

$$(4.6) \quad S(x, 0) = 80(1 - |1 - x|).$$

Here, we use nondimensional signal S . We suppose that initially the cells are uniformly distributed in the domain, at a cell density $n(x, 0) = n_0 \text{ mm}^{-1}$.

In Figure 4.1, we compare the stochastic simulation of the cell-based model with the solution of the macroscopic equations (4.4), (4.5) and (4.6). For the stochastic simulation, cell density is computed as the linear interpolation of the histogram for the positions of the cells. It is observed that the attractant concentration, cell density and cumulative cell density agree very well between the two models. We notice that in the first few minutes, an aggregate of cells forms because of the initial attractant gradient, later on the aggregate tends to be dispersed because diffusion smooths out the attractant gradient. We also notice that in this example $G'(S)\nabla S \cdot \mathbf{v}$ becomes as large as $30 \epsilon \text{ s}^{-1}$, but the solution of the chemotaxis equation (4.4) still provides a good approximation of the results of the cell-based model. This means that the chemotaxis equation may also be a good approximation of the underlying velocity jump process for a slightly weaker assumption than we used.

4.2. Self-organized aggregation in one dimensional space. In this example we investigate the motion of bacterial cells driven by the attractant that they produce. Thus the attractant dynamics is governed by

$$(4.7) \quad \frac{\partial S}{\partial t} = D_s \Delta S + \gamma n - \mu S.$$

We assume initially no attractant is added to the domain,

$$(4.8) \quad S(x, 0) = 0.$$

Periodic boundary conditions and the same parameters are used as in the first example. We set the initial condition of the cell density to be

$$(4.9) \quad n(x, 0) = n_0(1 + \text{small random noise}) \text{ mm}^{-1}$$

and stipulate that the noise has zero mean.

In Figure 4.2, we compare the stochastic simulation of the cell-based model with attractant dynamics (4.7, 4.8) and the solution of the continuum model (4.4, 4.9, 4.7, 4.8). We take $\mu = 1/3 \times 10^{-2} \text{ s}^{-1}$, $\gamma = 1/6 \times 10^{-1} / n_0 \text{ s}^{-1}$ per cell. A linear stability analysis (see Appendix B) of the continuum model around the uniform steady state (USS) $(n, S) \equiv (n_0, \gamma n_0 / \mu)$ shows that there are three unstable modes $\psi_k = e^{ik \frac{2\pi}{L} x}$, $k = 1, 2, 3$ with exponential growth rates 0.1439, 0.1954, 0.0904. Thus, we expect that instabilities develop around the uniform steady state and nonuniform peaks appear in the cell density profile. The system (4.4, 4.7) has no blow-up solutions in finite time [10], therefore nonuniform steady state develops finally.




FIG. 4.1. Aggregation and dispersion in a time-dependent signal field. The solution of equation (4.4) (smooth line) is compared with stochastic simulation of the cell based model when coupled with the attractant dynamics (4.5, 4.6). The left, center and right columns are the attractant concentration scaled by K_D , the cell density and cumulative cell density scaled by the average cell density n_0 at $t = 2, 5, 30$ and 90 min. $G(S)$, λ and $T(\mathbf{v}, \mathbf{v}')$ are given by equations (2.4, 4.3, 3.31). 4×10^3 cells are used for the Monte Carlo simulation ($n_0 = 10^3$). Other parameters used are $\lambda_0 = 1 \text{ s}^{-1}$, $b = 1 \text{ s}^{-1}$, $t_a = 2 \text{ s}$, $s = 20 \text{ } \mu\text{m/s}$, $K_D = 100$, $G_0 = 100$, $D_s = 8 \times 10^{-4} \text{ mm}^2/\text{s}$.




FIG. 4.2. Self-organized aggregation in bacterial colonies. (A)–(D): the solution of system (4.4, 4.9, 4.7, 4.8) is compared with one realization of the stochastic simulation of the cell based model coupled with the attractant dynamics given by equation (4.7, 4.8). The blue, green, red and cyan curves represent profiles taken at $t = 0, 20, 40, 180$ min. (E), (F): comparisons of the amplitudes of the first 4 Fourier modes of the solutions. Smooth lines: solution for the PDE system; dotted lines: stochastic simulation. 4×10^3 cells are used for the Monte Carlo simulation ($n_0 = 10^3$). $\mu = 1/3 \times 10^{-2} \text{ s}^{-1}$, $\gamma = 1/6 \times 10^{-4} \text{ s}^{-1}$ per cell. Other parameters used are the same as in Figure 4.1.

Figure 4.2 A – D show that in both models, the state of the system first evolves towards the unstable uniform steady state (green curve), then small perturbations finally lead the system to the stable nonuniform steady state (cyan curve). Because the perturbations in the two models are random and the periodic boundary condition allows for translation of solutions, we can not expect the peaks to appear at the same

x coordinate. Therefore neither averaging over different stochastic simulations of the cell-based model nor a point-wise comparison of the solutions of the two models is appropriate. Instead we compare the discrete Fourier coefficients ω_k of different modes $(\phi_k)_j = e^{k\frac{2\pi i}{N_x}j}$, $k, j = 0, 1, \dots, N_x - 1$ (Figure 4.2 E and F) in single realizations. We see that, in both models, the 0th mode amplitude ω_0 of n is constant because of the conservation of the total number of cells, and the 0th mode amplitude ω_0 of S increases first to the USS $\gamma n_0/\mu$ and stays at that value afterward. In the first few minutes, before reaching to the USS, the amplitude of the linearly-unstable modes ω_1 , ω_2 of n increases exponentially, and the amplitude of the other stable modes decreases exponentially (zoom-in of E). After the initial few minutes, due to the nonlinearity of the system, energy in the stable modes (both n and S) transfers to other modes, and coefficients ω_k increase until the system reaches the nonuniform steady state. We observed that in numerical calculations the exact time for the unstable modes to amplify sharply (around $t = 70 \text{ min} \sim 100 \text{ min}$ in this realization) depends strongly on the spectrum of the initial noise of the continuum model and the intrinsic noise of the cell-based model. Once the Fourier coefficients of the unstable modes exceeds a threshold (about 0.1 in this example), they start to grow faster than exponential. The amplitude of the most rapidly-varying modes of the cell-based model was observed to be much more noisy than that of the continuum model, because of the intrinsic time-dependent noise of the stochastic simulation.

To compare the two models in the case of multi-aggregate formation, we enlarge the domain from 4 mm to 8 mm to allow for more unstable modes. To match the number and location of the peaks in the early dynamics, we choose an initial cell density with sinusoidal noise

$$(4.10) \quad n(x, 0) = n_0(1 + \eta \sin(\frac{3\pi}{4}x) + \text{small random noise}).$$

In order to focus on the development of the instability, we set the signal at the uniform steady state initially

$$(4.11) \quad S(x, 0) = \frac{\gamma}{\mu}.$$

The numerical results for $\eta = 0.5$ are shown in Figure 4.3. We observe that aggregates form at the locations with maximum initial cell density (20 min, 40 min). Then, due to the instability of the multi-aggregate steady state, unevenness among different aggregates develops (180 min) and leads to merging of aggregates. Finally the single-aggregate stable steady state is reached (not shown). At $t = 20 \text{ min}$, the different noise in the two models are not significant, and the continuum model agrees well with the cell-based model (Figure 4.3 B). However, at $t = 40 \text{ min}$ and 180 min, the noise driven instability becomes important (Figure 4.3 C, D), and there one can not directly compare the exact value of the solution of the two models. However, it is shown that the two models exhibit the same qualitative dynamics in regard to merging of the aggregates.

From Figure 4.2, 4.3, we conclude that the dynamics of both models agree very well in different signal dynamics in 1-D, except for the location of the peaks where the signal gradient becomes $\mathcal{O}(1)$. In that case, as shown in section 3, the diffusion limit of the transport equation may not be a good approximation. In the next section, we apply both model in the background of network and aggregate formation in *E. coli* liquid assay.




FIG. 4.3. Multi-aggregate formation in bacterial colonies. In the top four plots, the time-elapse shots of the cell density solved from the continuum model (red line) is compared with one realization of the result of stochastic simulation of the cell based model (blue line) with initial conditions (4.10, 4.11). In the bottom two plots, the amplitude of the first 4 Fourier modes of the solutions are compared. Smooth lines: solution of the PDE system; dotted lines: stochastic simulation. 8×10^3 cells are used for the Monte Carlo simulation ($n_0 = 10^3$). Same parameters are used as in Figure 4.2.

4.3. Bacterial pattern formation: *E. coli* network and aggregate formation in liquid culture. When *E. coli* cells are suspended in a well-stirred liquid medium with succinate as the nutrient, they secrete the attractant aspartate and initially self-organize into a thread-like network, which quickly breaks into aggregates. The network-aggregate pattern appears on a time scale of 10 min. Since excess succinate is provided, cells grow in the exponential phase, and nutrient depletion is not

involved. In this example, we model the above dynamics in 2-D by both the hybrid cell-based approach and the macroscopic PDE approach, and compare the results.

The dynamics of the attractant is governed by the reaction-diffusion equation (4.7). The total cell number in the domain is N_0 and the average cell density n_0 . We use no-flux boundary conditions since there is no material exchange of the system with the environment. The uniform steady state of the continuum model (3.33, 4.7) is $(n, S) = (n_0, \gamma n_0/\mu)$. A linear analysis (see Appendix B) around the uniform steady state explains the pattern formation as the result of the amplification of the unstable modes of the fluctuation. To focus on the dynamics during pattern formation, we start from the uniform steady state with a small perturbation as the initial values,

$$(4.12) \quad n = n_0(1 + \text{small random noise}) \text{ mm}^{-2},$$

$$(4.13) \quad S(x, y, 0) = \gamma n_0/\mu.$$

In figure 4.4, We compare the numerical results of the continuum model (4.4, 4.7, 4.12, 4.13) with one realization of the stochastic simulation of the cell-based model. We used COMSOL Multiphysics to solve the 2-D continuum model (with 15648 triangles, using Lagrange elements), and the numerical algorithm given by Appendix A to simulate the cell-based model. The initial values for the continuum model are obtained by interpolating from the initial values of the cell-based model. Although the exact details of the transient dynamics can be different because of different noise in the two models, we note that both model predict comparable temporal and spatial features of the dynamical evolution from the network to the aggregates formation.

5. Chemotactic movement in external fields. Bacterial cells can swim in more complicated environments with external forces acting on them. For example, when the cell density becomes large, there may be mechanical interactions between cells, which may affect their swimming speed and direction. Another example arises when gravity becomes important. During the formation of bio-convection patterns reported in [11], oxygen-taxis drives the cells toward the top of the medium, while gravity acts downward. Therefore, the above analysis should finally be generalized to incorporate both forces between cells and forces due to external fields. The transport equation with external forces has the form

$$(5.1) \quad \frac{\partial p}{\partial t} + \nabla_x \cdot (\mathbf{v}p) + \nabla_v \cdot (\mathbf{a}p) + \nabla_y \cdot (\mathbf{f}p) = -\lambda(\mathbf{y})p + \int_V \lambda(\mathbf{y})T(\mathbf{v}, \mathbf{v}', \mathbf{y})p(\mathbf{x}, \mathbf{v}', \mathbf{y}, t) d\mathbf{v}'.$$

Previous results have been obtained for crawling cells [15], where the active force generation is incorporated by a simple cartoon description, and the jumps of velocity model random polarization of cells when no signal gradient is detected. Because the state inside each cell varies spatially, further dimension reduction is needed in that analysis.

Here we extend the analysis in section 3 to include external forces and consider a particular case in which bacteria swim close to a surface. In three dimensional space, bacterial cells swim in straight “runs”, but are subject to rotational diffusion. However, when they move near a surface, the “runs” display a consistent clockwise bias when observed from above [16, 12]. The bias can be explained by the interaction between the surface and the cell [19]. During a “run”, the cell body rotates clockwise




FIG. 4.4. *E. coli* network and aggregates formation. (A), (B): the cell density from the continuum model (A: $t = 7 \text{ min}$, B: $t = 13 \text{ min}$); (C), (D): the positions of the cells calculated from the cell-based model at the same time points; (E), (F): the interpolated cell density from (C) and (D). Parameters used include $\lambda_0 = 1 \text{ s}^{-1}$, $b = 5 \text{ s}^{-1}$, $t_a = 2 \text{ s}$, $s = 20 \text{ } \mu\text{m/s}$, $k_d = 40$, $D_s = 8 \times 10^{-4} \text{ mm}^2/\text{s}$, $\mu = 1/3 \times 10^{-2} \text{ s}^{-1}$, $\gamma = 1/6 \times 10^{-1}/n_0 \text{ s}^{-1}$, $n_0 = 400$.

while the flagella rotate counterclockwise when observed from behind. Therefore, when a cell swims parallel to a surface a larger viscous force is exerted on the bottom of the cell (closer to the surface) than that on the top of the cell, and thus net forces arise on both the cell body and the flagella, and these net forces induce the bias in the motion.

In the patterns formed in *P. mirabilis* colonies in [38], cells swim in a thin fluid-

like slime layer on top of the hard surface, and therefore the runs are biased. By incorporating a constant swimming bias to each cell's right, a two dimensional cell-based model leads to prediction of the chirality of spiral stream formation in *P. mirabilis* colonies [38]. In this section, we derive a corresponding macroscopic chemotaxis equation from the cell-based model with the swimming bias. We also incorporate persistence in the motion and thus assume the form of the turning kernel given by (3.34). The resulting equation enables us to see the interplay of chemotaxis and the swimming bias.

Let ω_0 be the constant angular velocity during a run. Then the acceleration has the form $\mathbf{a} = \omega_0 \mathbf{v} \times \mathbf{n}$, where \mathbf{n} is the normal vector of the surface pointing to the fluid side, *i.e.*, $\mathbf{a} = (\omega_0 v_2, -\omega_0 v_1)$. Let $p(\mathbf{x}, \mathbf{v}, z_2, t)$ be the cell density function. After nondimensionalization, the transport equation reads,

$$\begin{aligned} \epsilon^2 \frac{\partial p}{\partial t} + \epsilon \frac{\partial}{\partial x_1} (v_1 p) + \epsilon \frac{\partial}{\partial x_2} (v_2 p) + \omega_0 \frac{\partial}{\partial v_1} (v_2 p) - \omega_0 \frac{\partial}{\partial v_2} (v_1 p) \\ + \frac{\partial}{\partial z_2} \left(-\frac{z_2}{t_a} - G'(S) \left(\epsilon v_1 \frac{\partial S}{\partial x_1} + \epsilon v_2 \frac{\partial S}{\partial x_2} + \epsilon^2 \frac{\partial S}{\partial t} \right) p \right) \\ = (\lambda_0 + a_1 z_2 + a_2 z_2^2 + \dots) (-p + \int_V T(\mathbf{v}, \mathbf{v}') p(\mathbf{v}') d\mathbf{v}'). \end{aligned}$$

By multiplying 1, z_2^j/j , $j \geq 1$, and integrating with respect to z_2 , we get a system of equations for the z_2 -moments $\mathbf{M}(t, \mathbf{x}, \mathbf{v})$, where \mathbf{M} is defined as in (3.13),

$$\begin{aligned} (5.2) \quad \epsilon^2 \frac{\partial}{\partial t} \mathbf{\Lambda M} + \epsilon v_1 \frac{\partial}{\partial x_1} \mathbf{\Lambda M} + \epsilon v_2 \frac{\partial}{\partial x_2} \mathbf{\Lambda M} + \omega_0 v_2 \frac{\partial}{\partial v_1} \mathbf{\Lambda M} - \omega_0 v_1 \frac{\partial}{\partial v_2} \mathbf{\Lambda M} \\ = \epsilon^2 \mathbf{B M} + \epsilon \mathbf{C M} + \mathbf{D M}. \end{aligned}$$

If we apply the perturbation method directly to equation (5.2), there is no easy way to derive an approximating equation of the cell density, since M_0^0 is no longer independent of \mathbf{v} , and thus there is no simple relation between the cell density n and M_0^0 . Instead, we choose to proceed by multiplying (5.2) by 1, v_1 and v_2 , and integrating with respect to \mathbf{v} to get the complete moment equations. We define the density moments

$$n(\mathbf{x}, t) = \int M_0 d\mathbf{v}, \quad n_j(\mathbf{x}, t) = \int M_j d\mathbf{v}, \quad j = 1, 2, \dots, \quad \mathbf{n} = (n, n_1, n_2, \dots)^t,$$

and the velocity flux moments

$$\begin{aligned} J_{j,k}(\mathbf{x}, t) &= \int v_k M_j d\mathbf{v}, \quad j = 0, 1, 2, \dots, \quad \mathbf{J}_k = (J_{0,k}, J_{1,k}, J_{2,k}, \dots)^t, \quad k = 1, 2, \\ J_{j,kl}(\mathbf{x}, t) &= \int v_k v_l M_j d\mathbf{v}, \quad j = 0, 1, 2, \dots, \quad \mathbf{J}_{kl} = (J_{0,kl}, J_{1,kl}, J_{2,kl}, \dots)^t, \quad k, l = 1, 2. \end{aligned}$$

The subscript j is the index of the order of the z_2 -moment, and subscripts k, l are the indices of the velocity moment. We introduce the matrix operators

$$(5.3) \quad \mathbf{C}_k = -G'(S) \frac{\partial S}{\partial x_k} \text{diag}\{0, 1, 1, \dots\} \mathbf{J}^t, \quad k = 1, 2$$

where \mathbf{J} is the matrix operator defined in (3.18), but here acting on $l^\infty(\mathbb{R})$. We notice that componentwise $\mathbf{C}_1 + \mathbf{C}_2 = \mathbf{C}$ with \mathbf{C} defined by (3.16).

We also define matrix operators

$$(5.4) \quad \mathbf{D}_1 = -\text{diag} \left\{ 0, \frac{1}{t_a}, \frac{1}{t_a}, \frac{1}{t_a}, \dots \right\},$$

$$(5.5) \quad \mathbf{D}_2 = -\Lambda(\lambda_0 \mathbf{I} + \sum_{i=1}^{\infty} a_i \mathbf{J}_i)(1 - \psi_d) + \mathbf{D}_1.$$

To obtain the complete moment equations, we have to calculate $\int_V \mathbf{D} \mathbf{M} d\mathbf{v}$ and $\int_V v_k \mathbf{D} \mathbf{M} d\mathbf{v}$. Notice that, by property (ii) of \mathcal{A} , for any $f(\mathbf{v})$,

$$\int_V \mathcal{A} f d\mathbf{v} = \int_V \left(\int -I + T(\mathbf{v}, \mathbf{v}') d\mathbf{v} \right) f(\mathbf{v}') d\mathbf{v}' = 0,$$

therefore $\int_V \mathbf{D} \mathbf{M} d\mathbf{v} = \mathbf{D}_1 \mathbf{n}$. Assuming the turning kernel (3.34) and considering that

$$\int_V \mathbf{v} \mathcal{A} f d\mathbf{v} = \int_V \left(\int_V -\mathbf{v} f(\mathbf{v}) + \mathbf{v} T(\mathbf{v}, \mathbf{v}') d\mathbf{v} \right) f(\mathbf{v}') d\mathbf{v}' = -(1 - \psi_d) \int \mathbf{v} f(\mathbf{v}) d\mathbf{v},$$

we obtain

$$\int_V v_k \mathbf{D} \mathbf{M} d\mathbf{v} = \int_V \mathbf{D}_2 v'_k \mathbf{M}(\mathbf{v}') d\mathbf{v}' = \mathbf{D}_2 \mathbf{J}_k, \quad k = 1, 2.$$

Therefore the complete moment equations are

$$(5.6) \quad \epsilon^2 \frac{\partial}{\partial t} \Lambda \mathbf{n} + \epsilon \frac{\partial}{\partial x_1} \Lambda \mathbf{J}_1 + \epsilon \frac{\partial}{\partial x_2} \Lambda \mathbf{J}_2 = \epsilon^2 \mathbf{B} \mathbf{n} + \epsilon \mathbf{C}_1 \mathbf{J}_1 + \epsilon \mathbf{C}_2 \mathbf{J}_2 + \mathbf{D}_1 \mathbf{n},$$

$$(5.7) \quad \epsilon^2 \frac{\partial}{\partial t} \Lambda \mathbf{J}_1 + \epsilon \frac{\partial}{\partial x_1} \Lambda \mathbf{J}_{11} + \epsilon \frac{\partial}{\partial x_2} \Lambda \mathbf{J}_{12} - \omega_0 \Lambda \mathbf{J}_2 = \epsilon^2 \mathbf{B} \mathbf{J}_1 + \epsilon \mathbf{C}_1 \mathbf{J}_{11} + \epsilon \mathbf{C}_2 \mathbf{J}_{12} + \mathbf{D}_2 \mathbf{J}_1,$$

$$(5.8) \quad \epsilon^2 \frac{\partial}{\partial t} \Lambda \mathbf{J}_2 + \epsilon \frac{\partial}{\partial x_1} \Lambda \mathbf{J}_{12} + \epsilon \frac{\partial}{\partial x_2} \Lambda \mathbf{J}_{22} + \omega_0 \Lambda \mathbf{J}_1 = \epsilon^2 \mathbf{B} \mathbf{J}_2 + \epsilon \mathbf{C}_1 \mathbf{J}_{12} + \epsilon \mathbf{C}_2 \mathbf{J}_{22} + \mathbf{D}_2 \mathbf{J}_2.$$

Here \mathbf{B} is defined by (3.15). To close the moment equations, we follow [13] and assume the second velocity moments are isotropic, which is exact in 1-D:

$$(5.9) \quad J_{0,kl} = \frac{s^2}{2} n \delta_{kl}, \quad J_{j,kl} = \frac{s^2}{2} n_j \delta_{kl}, \quad k, l = 1, 2.$$

Then the moment equations reduce to

$$(5.10) \quad \epsilon^2 \frac{\partial}{\partial t} \Lambda \mathbf{n} + \epsilon \frac{\partial}{\partial x_1} \Lambda \mathbf{J}_1 + \epsilon \frac{\partial}{\partial x_2} \Lambda \mathbf{J}_2 = \epsilon^2 \mathbf{B} \mathbf{n} + \epsilon \mathbf{C}_1 \mathbf{J}_1 + \epsilon \mathbf{C}_2 \mathbf{J}_2 + \mathbf{D}_1 \mathbf{n},$$

$$(5.11) \quad \epsilon^2 \frac{\partial}{\partial t} \Lambda \mathbf{J}_1 + \epsilon \frac{\partial}{\partial x_1} \left(\frac{s^2}{2} \Lambda \mathbf{n} \right) - \omega_0 \Lambda \mathbf{J}_2 = \epsilon^2 \mathbf{B} \mathbf{J}_1 + \epsilon \mathbf{C}_1 \left(\frac{s^2}{2} \mathbf{n} \right) + \mathbf{D}_2 \mathbf{J}_1,$$

$$(5.12) \quad \epsilon^2 \frac{\partial}{\partial t} \Lambda \mathbf{J}_2 + \epsilon \frac{\partial}{\partial x_2} \left(\frac{s^2}{2} \Lambda \mathbf{n} \right) + \omega_0 \Lambda \mathbf{J}_1 = \epsilon^2 \mathbf{B} \mathbf{J}_2 + \epsilon \mathbf{C}_2 \left(\frac{s^2}{2} \mathbf{n} \right) + \mathbf{D}_2 \mathbf{J}_2.$$

Assuming the regular perturbation expansions, with superscript indicating the order of expansion,

$$\mathbf{n} = \mathbf{n}^0 + \epsilon \mathbf{n}^1 + \epsilon^2 \mathbf{n}^2 + \dots, \quad \mathbf{J}_k = \mathbf{J}_k^0 + \epsilon \mathbf{J}_k^1 + \epsilon^2 \mathbf{J}_k^2 + \dots, \quad k = 1, 2,$$

substituting into the moment equations (5.10-5.12), and comparing terms of equal orders of ϵ , we obtain,

$\mathcal{O}(\epsilon^0)$:

$$(5.13) \quad \mathbf{D}_1 \mathbf{n}^0 = 0,$$

$$(5.14) \quad \mathbf{D}_2 \mathbf{J}_1^0 = -\omega_0 \mathbf{\Lambda} \mathbf{J}_2^0,$$

$$(5.15) \quad \mathbf{D}_2 \mathbf{J}_2^0 = \omega_0 \mathbf{\Lambda} \mathbf{J}_1^0,$$

$\mathcal{O}(\epsilon^1)$:

$$(5.16) \quad \frac{\partial}{\partial x_1} \mathbf{\Lambda} \mathbf{J}_1^0 + \frac{\partial}{\partial x_2} \mathbf{\Lambda} \mathbf{J}_2^0 = \mathbf{C}_1 \mathbf{J}_1^0 + \mathbf{C}_2 \mathbf{J}_2^0 + \mathbf{D}_1 \mathbf{n}^1,$$

$$(5.17) \quad \frac{s^2}{2} \frac{\partial}{\partial x_1} \mathbf{\Lambda} \mathbf{n}^0 - \omega_0 \mathbf{\Lambda} \mathbf{J}_2^1 = \frac{s^2}{2} \mathbf{C}_1 \mathbf{n}^0 + \mathbf{D}_2 \mathbf{J}_1^1,$$

$$(5.18) \quad \frac{s^2}{2} \frac{\partial}{\partial x_2} \mathbf{\Lambda} \mathbf{n}^0 + \omega_0 \mathbf{\Lambda} \mathbf{J}_1^1 = \frac{s^2}{2} \mathbf{C}_2 \mathbf{n}^0 + \mathbf{D}_2 \mathbf{J}_2^1,$$

$\mathcal{O}(\epsilon^2)$:

$$(5.19) \quad \frac{\partial}{\partial t} \mathbf{\Lambda} \mathbf{n}^0 + \frac{\partial}{\partial x_1} \mathbf{\Lambda} \mathbf{J}_1^1 + \frac{\partial}{\partial x_2} \mathbf{\Lambda} \mathbf{J}_2^1 = \mathbf{B}_1 \mathbf{n}^0 + \mathbf{C}_1 \mathbf{J}_1^1 + \mathbf{C}_2 \mathbf{J}_2^1 + \mathbf{D}_1 \mathbf{n}^2.$$

From equation (5.13) we get $n_j^0 = 0$, $\forall j \geq 1$, or $\mathbf{n}^0 = (n^0, 0, 0, \dots)^t$. From equation (5.14, 5.15), we see that $(\mathbf{\Lambda}^{-1} \mathbf{D}_2)^2 \mathbf{J}_1^0 = -\omega_0^2 \mathbf{J}_1^0$, $(\mathbf{\Lambda}^{-1} \mathbf{D}_2)^2 \mathbf{J}_2^0 = -\omega_0^2 \mathbf{J}_2^0$. Since all the eigenvalues of $(\mathbf{\Lambda}^{-1} \mathbf{D}_2)^2$ are positive, it follows that $\mathbf{J}_1^0 = \mathbf{J}_2^0 = \mathbf{0}$.

Therefore equation (5.16) reduces to $\mathbf{D}_1 \mathbf{n}^1 = 0$, which means that $n_j^1 = 0$, $j \geq 1$, or $\mathbf{n}^1 = (n^1, 0, 0, \dots)^t$. Applying a similar argument to the 3rd and higher components of the equations (5.17, 5.18) gives $J_{j,1}^1 = J_{j,2}^1 = 0$, $\forall j \geq 2$. Thus the first two components of (5.17, 5.18) become

$$(5.20) \quad \frac{s^2}{2} \frac{\partial}{\partial x_1} n^0 - \omega_0 J_{0,2}^1 = -\lambda_0(1 - \psi_d) J_{0,1}^1 - a_1(1 - \psi_d) J_{1,1}^1,$$

$$(5.21) \quad -\omega_0 J_{1,2}^1 = -\frac{s^2}{2} G'(S) \frac{\partial S}{\partial x_1} n^0 - [\lambda_0(1 - \psi_d) + \frac{1}{t_a}] J_{1,1}^1,$$

$$(5.22) \quad \frac{s^2}{2} \frac{\partial}{\partial x_2} n^0 + \omega_0 J_{0,1}^1 = -\lambda_0(1 - \psi_d) J_{0,2}^1 - a_1(1 - \psi_d) J_{1,2}^1,$$

$$(5.23) \quad \omega_0 J_{1,1}^1 = -\frac{s^2}{2} G'(S) \frac{\partial S}{\partial x_2} n^0 - [\lambda_0(1 - \psi_d) + \frac{1}{t_a}] J_{1,2}^1.$$

From equations (5.21, 5.23), we find that

$$(5.24) \quad \begin{pmatrix} J_{1,1}^1 \\ J_{1,2}^1 \end{pmatrix} = -\frac{s^2 G'(S) n^0}{2(\lambda_0(1 - \psi_d) + \frac{1}{t_a})^2 + 2\omega_0^2} \begin{bmatrix} \lambda_0(1 - \psi_d) + \frac{1}{t_a} & \omega_0 \\ -\omega_0 & \lambda_0(1 - \psi_d) + \frac{1}{t_a} \end{bmatrix} \nabla S.$$

From equations (5.20, 5.22), we obtain

$$(5.25) \quad \begin{pmatrix} J_{0,1}^1 \\ J_{0,2}^1 \end{pmatrix} = -\frac{1}{\lambda_0^2(1 - \psi_d)^2 + \omega_0^2} \begin{bmatrix} \lambda_0(1 - \psi_d) & \omega_0 \\ -\omega_0 & \lambda_0(1 - \psi_d) \end{bmatrix} \cdot \left(\frac{s^2}{2} \nabla n^0 + a_1(1 - \psi_d) \begin{pmatrix} J_{1,1}^1 \\ J_{1,2}^1 \end{pmatrix} \right).$$

The first component of equation (5.19) is

$$(5.26) \quad \frac{\partial}{\partial t} n^0 + \frac{\partial}{\partial x_1} J_{0,1}^1 + \frac{\partial}{\partial x_2} J_{0,2}^1 = 0.$$

Substituting $J_{0,1}^1, J_{0,2}^1$ by equations (5.25) gives the final chemotaxis equation,

$$(5.27) \quad \frac{\partial}{\partial t} n^0 = D_n \Delta n^0 - \nabla \cdot [G'(S) n^0 (\chi_0 \nabla S + \beta_0 (\nabla S)^\perp)],$$

where

$$(5.28) \quad D_n = \frac{s^2}{2\lambda_0(1 - \psi_d) + \frac{2\omega_0^2}{\lambda_0(1 - \psi_d)}},$$

$$(5.29) \quad \chi_0 = \frac{a_1 s^2 (1 - \psi_d) [\lambda_0 (1 - \psi_d) (\lambda_0 (1 - \psi_d) + \frac{1}{t_a}) - \omega_0^2]}{2((\lambda_0 (1 - \psi_d) + \frac{1}{t_a})^2 + \omega_0^2) (\lambda_0^2 (1 - \psi_d)^2 + \omega_0^2)},$$

$$(5.30) \quad \beta_0 = \frac{\omega_0 a_1 s^2 (1 - \psi_d) (2\lambda_0 (1 - \psi_d) + \frac{1}{t_a})}{2((\lambda_0 (1 - \psi_d) + \frac{1}{t_a})^2 + \omega_0^2) (\lambda_0^2 (1 - \psi_d)^2 + \omega_0^2)},$$

and

$$(5.31) \quad \nabla S = \begin{pmatrix} \frac{\partial S}{\partial x_1} \\ \frac{\partial S}{\partial x_2} \end{pmatrix}, \quad (\nabla S)^\perp = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \nabla S.$$

From the forms of D_n , χ_0 and β_0 , we notice that when $\omega_0 = 0$, (5.27) reduces to the chemotaxis equation we derived in section 3.5 in a two-dimensional space. (5.27) can also be derived by using the assumptions in section 3.6. The macroscopic chemotactic velocity in (5.27) is given by

$$(5.32) \quad \mathbf{u}_S = G'(S) (\chi_0 \nabla S + \beta_0 (\nabla S)^\perp)$$

The magnitude of \mathbf{u}_S is

$$\begin{aligned} \|\mathbf{u}_S\| &= \|G'(S) \nabla S\| \sqrt{\chi_0^2 + \beta_0^2} \\ &= \|G'(S) \nabla S\| \cdot \frac{a_1 s^2 (1 - \psi_d)}{2\sqrt{((\lambda_0 (1 - \psi_d) + \frac{1}{t_a})^2 + \omega_0^2) (\lambda_0^2 (1 - \psi_d)^2 + \omega_0^2)}} \\ &= \|G'(S) \nabla S\| \cdot \frac{a_1 s^2 t_a}{2\lambda_0 (1 + (1 - \psi_d) \lambda_0 t_a)} \cdot \frac{1}{\sqrt{(1 + \frac{\omega_0^2}{(\lambda_0 (1 - \psi_d) + \frac{1}{t_a})^2}) (1 + \frac{\omega_0^2}{\lambda_0^2 (1 - \psi_d)^2})}}. \end{aligned} \quad (5.33)$$

The angle between \mathbf{u}_S and ∇S is

$$(5.34) \quad \theta_{\mathbf{u}_S, \nabla S} = \tan^{-1} \frac{\omega_0 (2\lambda_0 (1 - \psi_d) + \frac{1}{t_a})}{\lambda_0 (1 - \psi_d) (\lambda_0 (1 - \psi_d) + \frac{1}{t_a}) - \omega_0^2},$$

which is surprisingly independent of ∇S and a_1 .

5.1. Numerical comparison of the macroscopic chemotaxis velocity.

The analytical prediction of the macroscopic chemotaxis velocity (5.32) is shown to agree very well with statistics from the cell-based model at different signal gradients and bias levels ω_0 in Figure 5.1. Even for the large signal gradient $\|\nabla G(S)\| = 15$ (*i.e.*, $G'(S)\nabla S \cdot \mathbf{v} = 30 \text{ } \mu\text{s}^{-1}$), the difference is still within 10%.

The macroscopic chemotaxis velocity from the cell-based model is computed in the following way. For a given combination of $\nabla G(S)$ and ω_0 , we used $G(S) = S$, and a time-independent signal $S = Rx_2$ in order to guarantee $\nabla G(S)$ to be constant R in the whole path of a cell. Other parameters used remain the same as in previous examples. For each parameter combination, 6×10^3 cells are put at the same location with random initial velocity and zero initial y_2 . Positions of each cell are recorded every 1 min for a 30 min period. The position vector \mathbf{x}^i at time $t_i = i \text{ min}$ is computed by averaging all the cell positions. Then the macroscopic velocity vector is computed by applying the least square method to the averaged position, *i.e.*, by finding \mathbf{v} that minimizes $\sum_i (\mathbf{x}_j^i - \mathbf{v}t_i)$, where $j = 1, 2$ is the index for the space dimension.




FIG. 5.1. Comparison of the macroscopic velocity from equation (5.32, 5.34) with statistics from the cell-based model. In the first three plots, we compare $(u_S, v_S) = (\mathbf{u}_S \cdot \nabla S, \mathbf{u}_S \cdot (\nabla S)^\perp)$ as a function of $\nabla G(S)$ at different $\omega_0 = 0.02\pi, 0.04\pi, 0.06\pi$. Smooth lines are computed from the equation (5.32), dots are computed from the cell-based model; top lines and dots are for u_S , bottom ones are for v_S . The fourth plot is a comparison of the predicted angle $\theta_{\mathbf{u}_S, \nabla S}$ by equation (5.34) with simulation at different parameters. All other parameters are the same as the previous examples.

6. Chemotaxis induced by multiple signals. Single chemical induced chemotactic movement has been studied experimentally for various types of cells and modeled mathematically both microscopically and macroscopically [21, 37, 18]. However, many cell types are known to have multiple receptor types and thus can respond to many different chemicals. For instance, *E. coli* has five major types of receptors for

various nutrients, oxygen, etc. [37]. How these signals are integrated inside the cell is not generally known and may depend on the cell type. Macroscopic phenomenological chemotaxis equations have been proposed in [28]. In this section, we derive chemotaxis equations from a modified cell-based model by allowing multiple chemosignals.

In the case of *E. coli*, the signalling pathways for different chemicals share the same downstream phosphor-relaying network (including reactions of CheA, CheW, CheY, CheB, CheR, CheZ etc.), the only difference is the upstream transmembrane receptor. In the cell-based model in section 2, $G(S)$ describes detection of the signal, and \mathbf{y} describes the state of proteins within the cell. When there are multiple signals, G is generally a function of all possible signals, $G = G(S_1, S_2, \dots, S_m)$. By performing the standard procedure in section 3, a chemotaxis equation for multiple signals can be derived that has the following form

$$(6.1) \quad \frac{\partial}{\partial t} n = \nabla \cdot \left[D_n \nabla n - \chi_0 n \left(\frac{\partial G}{\partial S_1} \nabla S_1 + \dots + \frac{\partial G}{\partial S_m} \nabla S_m \right) \right],$$

where

$$(6.2) \quad \chi_0 = \frac{a_1 s^2 t_a}{N \lambda_0 (1 + (1 - \psi_d) t_a \lambda_0) (1 + (1 - \psi_d) t_e \lambda_0)}.$$

The functional form of G depends on the binding of the signal molecules to the receptors. Consider for example, the case of two attractants, and assume that all the binding is non-cooperative, and the two attractants S_1, S_2 competitively bind to the same receptor R as follows



Then according to the law of mass action, we have

$$\begin{aligned} \frac{d\overline{S_1}}{dt} &= -k_1^+ S_1 R + k_1^- \overline{S_1 R}, \\ \frac{d\overline{S_1 R}}{dt} &= +k_1^+ S_1 R - k_1^- \overline{S_1 R}, \\ \frac{d\overline{S_2}}{dt} &= -k_2^+ S_2 R + k_2^- \overline{S_2 R}, \\ \frac{d\overline{S_2 R}}{dt} &= +k_2^+ S_2 R - k_2^- \overline{S_2 R}, \\ \frac{dR}{dt} &= -k_1^+ S_1 R + k_1^- \overline{S_1 R} - k_2^+ S_2 R + k_2^- \overline{S_2 R}. \end{aligned}$$

If we further assume that the total number of receptors R_0 is conserved, then

$$R + \overline{S_1 R} + \overline{S_2 R} = R_0$$

Since the time scale of ligand binding is 10^{-2} s, which is small compared to the excitation and adaptation time, we may approximate the number of bound receptors by

the quasi-steady state value,

$$\begin{aligned}\overline{S_1 R} &= \frac{R_0 K_2 S_1}{K_1 K_2 + K_2 S_1 + K_1 S_2}, \\ \overline{S_2 R} &= \frac{R_0 K_1 S_2}{K_1 K_2 + K_2 S_1 + K_1 S_2},\end{aligned}$$

and G can be written as

$$G = g(\overline{S_1 R} + \overline{S_2 R}) = g\left(\frac{R_0(K_2 S_1 + K_1 S_2)}{K_1 K_2 + K_2 S_1 + K_1 S_2}\right).$$

If the two signals bind to different receptors, then a similar argument leads to the form,

$$G = g(\overline{S_1 R_1} + \overline{S_2 R_2}) = g\left(\frac{R_{10} S_1}{K_1 + S_1} + \frac{R_{20} S_2}{K_2 + S_2}\right).$$

In *E. coli*, the functioning units of chemoreceptors are observed to be trimers of dimers, and different types of receptors can form hetero-trimers *in vivo*; thus the form of the function of G actually can be even more complicated.

7. Discussion. In this paper, we developed a new method for deriving macroscopic equations of cell density from cell-level descriptions of chemotactic movement involved in bacterial pattern formation. The method involves solving the infinite \mathbf{y} -moment equations systematically by applying regular perturbation methods. It allows us to treat more general signal fields and cell-level descriptions than those used in the work [14, 13]. These generalizations include, allowing (a) time-dependent signal functions, (b) nonlinear turning frequency $\lambda(y_1)$ and (c) external force fields operating on cells. We also note that although we adopted the cartoon description of excitation and adaptation in their paper, the method proposed can manage autonomous ODE models with polynomial right-hand-sides, as long as the time scales of intracellular reactions can be separated from the diffusion scale of cells. Another advantage of the method is that it doesn't require unnecessary quasi-steady-state assumptions on the internal dynamics for closing the moments. The work is focused on studying the "run-and-tumble" chemotactic movement because it is the most understood biological system. However, the derivation method can also be applied to other systems without minor change. The limiting macroscopic equation maybe different depending on the details of the specific internal dynamics.

In the derivation of equation 3.33, we assumed that the signal detected by the cell in one "run" doesn't change significantly. This assumption is satisfied in most cases, but violation may appear inside self-organized aggregates. In this case, using a diffusion time and space scale may not be proper, and therefore a chemotaxis equation is not guaranteed to be a good approximation. However, in the numerical examples, we showed that the chemotaxis equation still captures the main dynamics of the cell-based model, although it seems to over-predict the amplitude and sharpness of the aggregates.

In the cell-based model, we assumed that cell density is low enough so that there is no mechanical interaction between cells. However, in some cases, the cell density can be high, and direct interaction between cells through the fluid, or more complicatedly, through their flagella can not be neglected. In these cases, we don't know so far how to incorporate these effects properly in the cell-based model, and if the microscopic




FIG. A.1. *Left: a schematic figure of the domains. The reaction-diffusion equations are solved on the grid, while the cells can move around the whole domain. Right: the area fractions used in defining the interpolators (A_1 , A_2).*

picture can be represented by macroscopic equations of cell density only. Phenomenological approaches have been proposed to incorporate the interaction as volume filling effects (need reference). However, how to justify their macroscopic equations in these cases need to be understood.

Appendix A. Numerical algorithm of the cell-based model. In the implementation of the cell-based model, cell motion is simulated by a standard Monte Carlo method in the whole domain, while the equations for extracellular chemicals are solved by an alternating direction method (Crank-Nicolson in 1-D) on a set of rectangular grid points (Figure A.1, left). In this appendix, we present the numerical algorithm in a two-dimensional domain with only one chemical – the attractant – involved (as in section 4.3). Each cell is described by its position (x_1^i, x_2^i) , internal variables (y_1^i, y_2^i) , direction of movement θ^i and age T^i (the superscript i is the index of the cell). Concentration of the attractant is described by a discrete function defined on the grid for the finite difference method (Figure A.1, left). We denote the time step by k , the space steps by h_1 and h_2 . Since two components of the model live in different spaces, two interpolating operators are needed in the algorithm. \mathcal{T}_{gc} is used to evaluate the attractant concentration that a cell senses. For a cell at (x_1^i, x_2^i) , inside the square with vertex indices $(n-1, m-1)$, $(n, m-1)$, $(n-1, m)$ and (n, m) , $\mathcal{T}_{gc}(x_1^i, x_2^i)$ is defined by the bi-linear function:

$$(A.1) \quad \mathcal{T}_{gc}(x_1^i, x_2^i) = \frac{A_4}{A} S_{n-1, m-1} + \frac{A_3}{A} S_{n, m-1} + \frac{A_2}{A} S_{n-1, m} + \frac{A_1}{A} S_{n, m}$$

where $A = h_1 h_2$ and $A_j, j = 1, 2, 3, 4$ are the area fractions (Figure A.1, right). On the other hand, the attractant secreted by cells is interpolated as increments at the grid points by \mathcal{T}_{cg} . Suppose during one time step k , a cell staying at (x_1^i, x_2^i) secretes Δ amount of attractant, we then interpolate the increment of the attractant

concentration at the neighboring grid points as follows:

$$(A.2) \quad \mathcal{T}_{cg}(\mathbf{x}^i; p, q) = \begin{cases} \frac{A_4 \Delta}{A^2}, & (p, q) = (n-1, m-1); \\ \frac{A_3 \Delta}{A^2}, & (p, q) = (n, m-1); \\ \frac{A_2 \Delta}{A^2}, & (p, q) = (n-1, m); \\ \frac{A_1 \Delta}{A^2}, & (p, q) = (n, m); \\ 0, & \text{otherwise.} \end{cases}$$

We consider here a periodic boundary condition. The detailed computing procedure is summarized as follows.

S1 Initialization.

- (a) Initialize the chemical fields.
- (b) Initialize the list of swimmer cells. Each cell is put in the domain with random position, moving direction and age. \mathbf{y}^i is set to be 0.

S2 For time step l ($= 1$ initially), update the data of each cell.

- (a) Determine the direction of movement θ^i by equation (3.34).
 - i) Generate a random number $r \in U[0, 1]$;
 - ii) If $r < 1 - e^{-\lambda^i k}$, update θ^i with a new random direction.
- (b) $(x_1^i, x_2^i)_l \leftarrow (x_1^i, x_2^i)_{l-1} + (sk \cos \theta^i, sk \sin \theta^i)$. Apply periodic boundary condition to make sure (x_1^i, x_2^i) inside the domain,
- (c) $(T^i)_l \leftarrow (T^i)_{l-1} + k$. If $(T^i)_l \geq 2$ hours, then divide the cell into two daughter cells. This step is only considered when cell growth is considered.
- (d) Update (y_1^i, y_2^i) by equations (4.1, 4.2).
 - i) Determine the attractant concentration before the cell moves $(S^i)_{l-1}$ and after the cell moves $(S^i)_l$ by using the interpolating operator \mathcal{T}_{gc} .
 - ii) Estimate the attractant level during the movement by $S^i(t) = (S^i)_{l-1} \frac{t-lk}{k} + (S^i)_l \frac{lk+k-t}{k}$ and integrate equation (4.1) to get $(y_2^i)_l$.
 - iii) $(y_1^i)_l \leftarrow G(S) - (y_2^i)_l$.

S3 Compute the source term of the attractant $f^{l-\frac{1}{2}}$ due to the secretion by the cells using the interpolator \mathcal{T}_{cg}

$$f_{p,q}^{l-\frac{1}{2}} = \sum_i (\mathcal{T}_{cg}((\mathbf{x}^i)_{l-\frac{1}{2}}; p, q)),$$

where $\Delta = \gamma k$.

S4 Apply the alternating direction implicit method to the equation of the attractant (4.5):

$$\begin{aligned} \frac{S_{p,q}^{l-1/2} - S_{p,q}^{l-1}}{k/2} &= D_s \frac{S_{p+1,q}^{l-1/2} - 2S_{p,q}^{l-1/2} + S_{p-1,q}^{l-1/2}}{h_x^2} \\ &\quad + D_s \frac{S_{p,q+1}^{l-1} - 2S_{p,q}^{l-1} + S_{p,q-1}^{l-1}}{h_y^2} - \gamma \frac{S_{p,q}^{l-1} + S_{p,q}^{l-1/2}}{2} + f_{p,q}^{l-\frac{1}{2}}, \\ \frac{S_{p,q}^l - S_{p,q}^{l-1/2}}{k/2} &= D_s \frac{S_{p+1,q}^{l-1/2} - 2S_{p,q}^{l-1/2} + S_{p-1,q}^{l-1/2}}{h_x^2} \\ &\quad + D_s \frac{S_{p,q+1}^l - 2S_{p,q}^l + S_{p,q-1}^l}{h_y^2} - \gamma \frac{S_{p,q}^{l-1/2} + S_{p,q}^l}{2} + f_{p,q}^{l-\frac{1}{2}}. \end{aligned}$$

For the boundary grid points, use the periodic scheme.

S5 $l \leftarrow l + 1$. If $lk \leq T_0$, repeat **S2-S4**; otherwise, return.

Appendix B. linear analysis on the stability of the uniform steady state of the continuum model.

Linear analysis of the classical chemotaxis equation system has been done in the literature (need reference). For readers' convenience, we include it here. Consider the system

$$(B.1) \quad \frac{\partial n}{\partial t} = \nabla \cdot (D_n \nabla n - \chi(S)n \nabla S)$$

$$(B.2) \quad \frac{\partial S}{\partial t} = D_s \Delta S + \gamma n - \mu S$$

The uniform steady state is $(n, S) = (n_0, \frac{\gamma}{\mu} n_0)$, where n_0 is the averaged cell density. Without loss of generality, we assume a one-dimensional domain $[0, L]$ with periodic boundary conditions, as in the example 4.1 and 4.2. The analysis can be extended without difficulty in the two dimensional case of example 4.3.

Let $u = n - n_0$, $v = S - S_0$ with $S_0 = \frac{\gamma}{\mu} n_0$. By linearizing around the uniform steady state, we get the system

$$(B.3) \quad \frac{\partial u}{\partial t} = D_n \Delta u - \chi(S_0)n_0 \Delta v$$

$$(B.4) \quad \frac{\partial v}{\partial t} = D_s \Delta v + \gamma u - \mu v$$

Assume

$$u = \sum_{q \neq 0} \varphi_q(t) e^{i\omega_q x}, \quad v = \sum_{q \neq 0} \psi_q(t) e^{i\omega_q x}$$

with $\omega_q = 2q\pi/L$. The system can thus be reduced to

$$(B.5) \quad \begin{pmatrix} \varphi_q \\ \psi_q \end{pmatrix}_t = A_q \begin{pmatrix} \varphi_q \\ \psi_q \end{pmatrix} \quad \text{with} \quad A_q = \begin{pmatrix} -D_n \omega_q^2 & \chi(S_0)n_0 \omega_q^2 \\ \gamma & -D_s \omega_q^2 - \mu \end{pmatrix}$$

for any $q \in \mathbb{Z}, q \neq 0$. The uniform steady state of the nonlinear system is unstable if the linearized system (B.3 - B.4) has exponentially growing non-homogeneous modes, which means that there exists a wave number q such that A_q has a positive eigenvalue.

Simple calculation leads to

- $Tr(A_q) = -(D_n + D_s)\omega_q^2 - \mu < 0$
- $Det(A_q) = \omega_q^2(D_n D_s \omega_q^2 + \mu D_n - \gamma \chi(S_0)n_0)$
- $Tr(A_q)^2 - 4Det(A_q) = [(D_s - D_n)\omega_q^2 + \mu]^2 + 4\gamma \omega_q^2 \chi(S_0)n_0 > 0$.

Therefore A_q is simple with eigenvalues

$$(B.6) \quad \lambda_q^\pm = \frac{Tr(A_q) \pm \sqrt{Tr(A_q)^2 - 4Det(A_q)}}{2}$$

λ_q^- is always negative and approaches $-\infty$ as $n, m \rightarrow \infty$, λ_q^+ can be positive when $Det(A_q) = \lambda_q^- \lambda_q^+ < 0$. Thus, the instability condition of the uniform steady state is,

$$\exists q \neq 0, \text{ s.t. } Det(A_q) < 0 \Leftrightarrow \omega_q^2(D_n D_s \omega_q^2 + \mu D_n - \gamma \chi(S_0)n_0) < 0$$

which is equivalent to

$$(B.7) \quad 4\pi^2 D_s + L^2 \left(\mu - \frac{\gamma \chi (\gamma / \mu n_0) n_0}{D_n} \right) < 0$$

The growth rate of an unstable mode in the linear system is given by λ_q^+ .

From the instability analysis, we can also see that if the q th mode is unstable, the lower modes are always unstable. This leads to multiple nonuniform steady states of the nonlinear system (B.1, B.2), but only the one with a single high peak is stable.

REFERENCES

- [1] J. ADLER, *Chemotaxis in bacteria*, SCIENCE, 153 (1966), pp. 708–716.
- [2] E. BEN-JACOB, I. COHEN, O. SHOCHET, I. ARANSON, AND H. LEVINE, *Complex bacterial patterns*, Nature, 373 (1995), pp. 566–557.
- [3] H. C. BERG, <http://webmac.rowland.org/labs/bacteria/movies/others/>.
- [4] ———, *Random Walks in Biology*, Princeton University Press, 41 William Street, Princeton, New Jersey 08540, 1983.
- [5] ———, *Motile behavior of bacteria*, Physics Today, 53 (2000), pp. 24–29.
- [6] H. C. BERG AND D. BROWN, *Chemotaxis in Escherichia Coli analyzed by three-dimensional tracking.*, Nature, 239 (1972), pp. 502–507.
- [7] R. BOURRET, BORKOVICH K, AND M. SIMON, *Signal transduction pathways involving protein phosphorylation in prokaryotes*, Annu. Rev. Biochemistry, 60 (1991), pp. 401–441.
- [8] M. BRENNER, L. LEVITOV, AND E. BUDRENE, *Physical mechanisms for chemotactic pattern formation by bacteria*, Biophys J., 74 (1998), pp. 1677–93.
- [9] E. O. BUDRENE AND H. C. BERG, *Dynamics of formation of symmetrical patterns by chemotactic bacteria.*, Nature, 376 (1995), pp. 49–53.
- [10] S. CHILDRESS AND J. K. PERCUS, *Nonlinear aspects of chemotaxis*, Mathematical Biosciences, 56 (1981), pp. 217–237.
- [11] A. CZIRÓK, I. M. JÁNOSI, AND J. O. KESSLER, *Bioconvective dynamics: dependence on organism behaviour.*, J. Exp. Biol., 203 (2000), pp. 3345–54.
- [12] W. R. DILUZIO, L. TURNER, M. MAYER, P. GARSTECKI, D. B. WEIBEL, H. C. BERG, AND G. M. WHITESIDES, *Escherichia coli swim on the right-hand side*, Nature, 435 (2005), pp. 1271–4.
- [13] R. ERBAN AND H. OTHMER, *From signal transduction to spatial pattern formation in e. coli: A paradigm for multiscale modeling in biology*, Multiscale Model. Simul., 3 (2005), pp. 362–394.
- [14] R. ERBAN AND H. G. OTHMER, *From individual to collective behavior in bacterial chemotaxis*, SIAM J. Appl. Math., 65 (2004), pp. 361–391.
- [15] ———, *Taxis equations for amoeboid cells*, J Math Biol, 54 (2007), pp. 847–885.
- [16] P. D. FRYMIER, R. M. FORD, H. C. BERG, AND P. T. CUMMINGS, *Three-dimensional tracking of motile bacteria near a solid planar surface*, PNAS, 92 (1995), pp. 6195–6199.
- [17] T. HILLEN AND H. G. OTHMER, *The diffusion limit of transport equations derived from velocity-jump processes*, SIAM J. Appl. Math., 61 (2000), pp. 751–775.
- [18] D. HORSTMANN, *From 1970 until present: the Keller-Segel model in chemotaxis and its consequences I*, Jahresbericht der DMV, 105 (2003), pp. 103–165.
- [19] E. LAUGA, W. R. DILUZIO, G. M. WHITESIDES, AND H. A. STONE, *Swimming in circles: Motion of bacteria near solid boundaries*, Biophys. J., 90 (2006), pp. 400–412.
- [20] J. T. LOCSEI, *Persistence of direction increases the drift velocity of run and tumble chemotaxis*, J Math Biol, 55 (2007), pp. 41–60.
- [21] R. LUX AND W. SHI, *Chemotaxis-guided movements in bacteria*, Crit. Rev. Oral. Biol. Med., 15 (2004), pp. 207–20.
- [22] R. M. MACNAB, *Sensing the environment: Bacterial chemotaxis*, in Biological Regulation and Development, R. Goldberg, ed., New York, 1980, Plenum Press, pp. 377–412.
- [23] B. A. MELLO AND Y. TU, *Quantitative modeling of sensitivity in bacterial chemotaxis: the role of coupling among different chemoreceptor species*, Proc. Nat. Acad. Sci. (USA), 100 (2003), pp. 8223–8228.
- [24] N. MITTAL, E. O. BUDRENE, M. P. BRENNER, AND A. OUDENAARDEN, *Motility of escherichia coli cells in clusters formed by chemotactic aggregation*, Proc Natl Acad Sci U S A, 100 (2003), pp. 13259–63.

- [25] H. G. OTHMER, S. R. DUNBAR, AND W. ALT, *Models of dispersal in biological systems.*, J. Math. Biol., 26 (1988), pp. 263–298.
- [26] H. G. OTHMER AND T. HILLEN, *The diffusion limit of transport equations ii: chemotaxis equations*, SIAM J. Appl. Math., 62 (2002), pp. 1222–1250.
- [27] H. G. OTHMER AND P. SCHAAP, *Oscillatory cAMP signaling in the development of Dictyostelium discoideum*, Comments on Theoretical Biology, 5 (1998), pp. 175–282.
- [28] K. J. PAINTER, P. K. MAINI, AND H. G. OTHMER, *Development and applications of a model for cellular response to multiple chemotactic cues*, J. Math. Biol., 41 (2000), pp. 285–314.
- [29] GEORGE C. PAPANICOLAOU, *Asymptotic analysis of transport processes*, Bulletin of the American Mathematical Society, 81 (1975), pp. 330–393.
- [30] A. A. POLEZHAEV, R. A. PASHKOV, A. I. LOBANOV, AND I. B. PETROV, *Spatial patterns formed by chemotactic bacteria Escherichia coli*, Int. j. dev. biol., 50 (2006), pp. 311–338.
- [31] C. V. RAO, J. R. KIRBY, AND A. P. ARKIN, *Design and diversity in bacterial chemotaxis: A comparative study in Escherichia coli and bacillus subtilis.*, PLoS Biol, 2 (2004), p. E49.
- [32] T. S. SHIMIZU, S. V. AKSENOV, AND D. BRAY, *A spatially extended stochastic model of the bacterial chemotaxis signalling pathway.*, J Mol Biol., 329 (2003), pp. 291–309.
- [33] P. A. SPIRO, J. S. PARKINSON, AND H. G. OTHMER, *A model of excitation and adaptation in bacterial chemotaxis*, Proc. Natl. Acad. Sci. USA, 94 (1997), pp. 7263–7268.
- [34] L. TSIMRING, H. LEVINE, I. A., E. BEN-JACOB, I. COHEN, O. SHOCHET, AND W. N. REYNOLDS, *Aggregation patterns in stressed bacteria*, 75 (1995), pp. 1859–1862.
- [35] R. TYSON, S. R. LUBKIN, AND J. D. MURRAY, *A minimal mechanism for formation*, Proc. R. Soc. Lond. B, 266 (1999), pp. 299–304.
- [36] ———, *Model and analysis of chemotactic bacterial patterns in a liquid medium*, J. Math. Biol., 38 (1999), pp. 359–375.
- [37] G. WADHAMS AND J. ARMITAGE, *Making sense of it all: bacterial chemotaxis*, Nat. Rev. Mol. Cell Biol., 5 (2004), pp. 1024–37.
- [38] C. XUE, E. O. BUDRENE-KAC, AND H. G. OTHMER, *Radial and spiral streams formation in bacterium Proteus mirabilis colonies*, preprint, (2007).